nss e n

## Conclusion

COVID-19 **\( \mathbb{C}** \) \( \mathbb{C} \) \( \mathbb{M} \) \( \mathbb{C} \) \( \mathbb{M} \) \( \mathbb{C} \) \( \mathbb{C

- with thoracic life-limiting illnesses: lung cancers and non-malignant end-stage respiratory diseases. Thorax 76: 989-995.
- Bourke SJ, Peel ET (2014) Palliative care of chronic progressive lung disease. Clin Med 14: 79-82.
- Hyasat K, Sriram KB (2016) Evaluation of the patterns of care provided to patients With COPD compared to patients with lung cancer who died in hospital. Am J Hosp Palliat Care 33: 717-722.
- 8. Gutierrez Sanchez D, Perez Cruzado D, Cuesta-Vargas AI (2018) The quality of dying and death measurement instruments: A systematic psychometric review. J Adv Nurs 74: 1803-1808.
- Oh YM, Kang YN, Han SJ, Kim JH (2023) Decision and Practice of End-of-Life Care in Lung Disease Patients with Physicians Orders for Life Sustaining Treatment. Korean J Hosp Palliat Care 26: 7-17.
- Lee MA (2019) Withdrawal of life-prolonging medical care and hospicepalliative care. J Korean Med Assoc 62: 369-375.

\$