

Achilles tendon; Spinal anesthesia; Percutaneous repair

The treatment modalities for Achilles tendon rupture vary from conservative to different surgical treatment options. The main concerns of conservative treatment are the complications of long term immobilization and re-rupture rate, which could be minimized by advances in postoperative rehabilitation programs and functional bracing. On the contrary, the main concern of open repair is wound complications, which could be minimized by recent advances in minimally invasive techniques. Percutaneous repair of acute tendoachilles rupture avoids possible complications of open repair as well as conservative treatment [1]. This paper presents a series of patients with acute tendon rupture treated with percutaneous repair carried. The purpose of this study is to evaluate the clinical and functional results of percutaneous repair of acute Achilles tendon ruptures and efficacy of this technique in sural nerve protection.

The study was approved by the ethical committee of Dpu University, Pune. A total of 22 patients with complete rupture of the Achilles tendon were treated in Dr. D. Y. Patil hospital and medical college pune, using the presented technique. All patients signed an informative consent form.

There were 16 men and 6 women, with a mean age of 34.7 years (25 years-48 years). The causes of injury were sports activity (12 patients) twist injury (six), falling downstairs (two) and during daily activity (two).

The rupture was diagnosed at the site of rupture a palpable tendon gap, positive Thompson test, positive matles test and loss of plantar flexion in the affected ankle and unable to walk on toes on the affected side (Figure 1). The diagnosis was confirmed by radiological evaluation (MRI/USG) for all patients (Figure 2). All patients had a complete rupture of the Achilles tendon with a gap less than 3 cm located 3 cm-6 cm proximal to the calcaneal insertion[2]. The inclusion criteria were complete rupture of Achilles tendon of not more than 1 month duration with intact skin and intact sensation of the affected limb. Exclusion criteria were as follows (a) incomplete rupture of Achilles tendon, (b) chronic rupture exceeding 1 month duration, (c) sensory impairment of the affected limb, (d) bony avulsion of calcaneal tuberosity, (e) history of recent local corticosteroid injection and (f) re-rupture of the Achilles' tendon. The percutaneous tendon repair was done with in a mean of 8.5 days (2 days-28 days) of injury. The mean follow up was 26 months (18 months-40 months).

**\*Corresponding author:** Purvam P. Jivrajani, Department of Orthopedics, Dr. D.Y. Patil Medical College and hospital and research centre, Pune, E-mail: purvamjivrajani@gmail.com

**Received:** July 02, 2021; **Accepted:** July 16, 2021; **Published:** July 23, 2021

**Citation:** Jivrajani PP, Biswas SK, Saikishan S, Anurag A, Shubham T, et al. (2021) Percutaneous Repair of Achilles Tendon Rupture. Clin Res Foot Ankle 9: 314.

**Copyright:**

All patients underwent surgery under spinal anesthesia without tourniquet. Patients were placed in prone position with free ankle and foot for easily mobilization [3]. After detecting the site of rupture, skin incision markings were marked along the posterior aspect of the Achilles' tendon. Leg and foot were prepared and draped.

- a. Needles 16G ss needless short and long
- b. Sutures ethibond (5-0), ethilon (3-0) (Figure 3)

### 5: Schematic Diagrams of Procedure

#### 3: Ethibond can be easily passed through the 16G needle

Over the posterior aspect of tendon, 4 cm-5 cm above the proximal ruptured end of tendon, a 16G needle is passed from medial to lateral through the tendon and ethibond is passed through the 16G needle. Then two 16G needles are passed obliquely inferior to superior through the tendon and ethibond is passed through the needles (Figures 4-6).

#### 6: 16G needles are passed obliquely inferior to superior through the tendon

This similar technique is followed again till the proximal end of tendon is held firmly with ethibond. Then this similar technique is followed in the distal part of Ruptured tendon. Keep the ankle in plantar flexion till you get approximation of ruptured ends of tendon [4]. Keeping the ankle in plantar flexion, drill bit (2.7 mm) is passed through the calcaneum medial to lateral, and 16G needle is passed through the drill hole and lateral end of ethibond is passed through the needle (laterally to medially) so that both the free ends of ethibond are placed medially, free ends are tied and knot is buried underneath the skin, if necessary apply skin sutures. Hold the limb in plantar flexion and apply pop cast after aseptic dressing of wound, pop should be moulded along the course of tendoachilles (Figures 7-10).

### 7: Schematic Diagrams of Procedure

#### 4: Operative Technique

