Hersini, et al., J Pain Relief 2015, 4:5
DOI: 10.4172/2167-0846.1000201

Research Article Open Access

Median pain intensity of acute pain was 5 (IQR: 4.75) and mean current pain was 2.8 (SD: 2.4) on the NRS.

Twenty-one patients (55%) took medication for chronic pain including paracetamol (n=20), opioids (n=17), NSAIDs (n=4), gabapentin (n=1), and pregabalin (n=1). One patient reported using physiotherapy in addition to pain killers for pain management.

The McGill Pain Questionnaire was used to explore the patients´ subjective experience of their pain. All PTPS patients completed the questionnaire. The mean number of descriptive words chosen was 6.9 (SD: 5.2). The most common terms selected to describe the pain were sensory words, e.g. shooting, crushing, and aching. The three most frequently descriptors used were "annoying" (18/37), "shooting" (14/37), and "stabbing" (14/37).

Neuropathic pain symptoms

All PTPS patients completed the questionnaire. Of the 38 patients with PTPS, 36 (95%) experienced neuropathic pain characteristics (NPSI score>0) [11]. The intensity and proportion of patients reporting each neuropathic pain symptom are shown in Table 3. Pressing (deep) pain (symptoms of squeezing and pressing) was the most common reported term (69%), followed by evoked pain (pain evoked by pressure, brush or cold) (61%). Paresthesia/dysesthesia (tingling and pins and needles) was seen in 56% of patients, and paroxysmal pain (symptoms of stabbing and electric shocks) in 53%. Burning (superficial) pain was seen in 22% of the patients with PTPS.

Predictors of post-thoracotomy pain syndrome

Acute post-operative pain measured by VAS was significantly higher in patients with PTPS than those without chronic pain ($p< 0.01$). Furthermore, patients with PTPS had a longer hospital stay than patients with no chronic pain ($p=0.05$).

There were no significant differences between patients with and without PTPS concerning gender or age at the study, age at the surgery, time since the surgery, operation side, operational procedure, extent of surgery, postoperative outcome, pre- or postoperative radiotherapy/chemotherapy, TNM stage, or preoperative pain (Table 4).

Quality of life

Quality of life was measured using the SF-36 questionnaire. The questionnaire was completed by 131 patients (98.5%). The mean total score was 47 (SD: 21) for patients with PTPS and 69 (SD: 21.5) for

	PTPS (n=38)	No pain (n=95)	p-value
Sex: Male	15 (39 %)	53 (56%)	0.13
Operation side: Left	19 (50%)	43 (45%)	0.76
Operational procedure			0.65
VATS	9 (24%)	28 (29%)	
Antero- or posterolateral thoracotomy	29 (76%)	67 (71 %)	
Age at study (Years), Mean (SD)	63.9 (10.6)	67.4 (9.9)	0.07
Age at surgery (years), Mean (SD)	61.9 (10.7)	65.3 (9.8)	0.10
Time since surgery (months), Mean (SD)	23 (12)	27.2 (11.9)	0.07
Extent of surgery			0.56
Wedge resection	1 (3%)	7 (8%)	
Lobectomy	27 (71%)	71 (75%)	
Lobectomy with wedge-resection	2 (5%)	4 (4%)	
Lobectomy and resection (other)	3 (8%)	2 (2%)	
Bilobectomy	2 (5%)	6 (6%)	
Pneumonectomy	3 (8%)	5 (5%)	
Resection of ribs or latissimus dorsi muscles (yes/no)	1(3%)/37(97%)	3(3%)/93(97%)	1
Postoperative outcome			0.63
Pneumothorax/Hydrothorax/Hydropneumothorax	5 (13%)	10 (11%)	
Pyothorax	1 (3%)	1 (1%)	
Subcutaneous emphysema	2 (5%)	1 (1%)	
Atelectasis	2 (5%)	3 (3%)	
Other	2 (5%)	5 (5%)	
TNM			0.33
T1	18 (47%)	41 (43%)	
T2	14 (37%)	35 (37%)	
T3	4 (11%)	18 (19%)	
T4	2 (5 %)	1 (1%)	
Acute pain (VAS after operation), Median (IQR)	5 (4.8)	3 (5)	<0.01
Preoperative Radiotherapy/Chemotherapy (yes/no)	2(5%)/36(95%)	10(11%)/85(89%)	0.53
Postoperative Radiotherapy/Chemotherapy (yes/no)	9(24%)/29(76%)	30(32%)/65(68%)	0.49
Length of hospital stay (days), Median (IQR)	10 (7)	8 (4)	<0.05
Preoperative pain (yes/no)	3(8%)/35(92%)	11(12%)/84(88%)	0.75

Abbreviations: PTPS: Post-Thoracotomy Pain Syndrome; VATS: Video-Assisted Thoracoscopic Surgery; SD: Standard Deviation; IQR: Interquartile Range; VAS: Visual Analogue Scale. Data are presented in numbers of patients unless stated otherwise

Table 4: Predictive factors for the prevalence of post-thoracotomy pain syndrome.

SF-36 scales	PTPS (n=93)	No pain (n=38)	p-value
Physical functioning	47.4 (21.7)	69.2 (25.4)	<0.01
Role limitations – physical problems	20.4 (32.3)	54.0 (41.9)	<0.01
Bodily pain	52.0 (25.4)	81.0 (26.1)	<0.01
General health perception	44.4 (21.0)	63.2 (21.8)	<0.01
Vitality	40.5 (21.1)	[illegible]	[illegible]

Preoperative pain has been shown to be a risk factor for persistent pain in several types of surgery [19-21], but only few studies on PTPS has assessed preoperative pain and these have shown conflicting results [14,22,23]. The role of preoperative pain in the development of chronic pain following thoracic surgery should be considered in future studies.

Post-operative use of analgesics

Half of the patients with PTPS used analgesics for their pain following surgery. Surprisingly, 81% of the patients using analgesics took opioids (17/21), which are normally used for moderate to severe pain. This could indicate that the pain is disabling although reported as mild. Only 9.5% (2/21) used antineuropathic medication (gabapentin and pregabalin) in contrast to 95% with a neuropathic component according to NPSI. This is in accordance with a previous study which found that only 2% used antineuropathic medication in contrast to 23% with a neuropathic component according to the PainDETECT Questionnaire [4].

Study limitations

The design of the current study holds limitations that should be kept in mind. The current study estimated the prevalence of pain based on a self-reported pain by patients, which can be a potential bias to the results due to recall bias. Furthermore, the questionnaire constructed by the first author has never been validated on its own. However, questions asked were simple and we have no reason to believe that answers differ between surgical groups. Objective assessment of pain could support the self-reported data. The cross-sectional design of this study provides a point prevalence of PTPS estimated within a specific time point, and, therefore, some patients might not have developed pain yet, and some patients could have had chronic pain after surgery that have disappeared at the time of the study. In addition, the population of the study may not have been enough large to proof statistical significance.

Non-responders are another source of bias. If the prevalence of PTPS among the non-responders was high, this study might have underestimated the prevalence of PTPS and vice versa. The non-responders could also have different characteristics of pain thaonds0

18. Grosen K, Laue Petersen G, Pfeiffer-Jensen M, Hoejsgaard A, Pilegaard HK (2013) Persistent post-surgical pain following anterior thoracotomy for lung cancer: a cross-sectional study of prevalence, characteristics and interference with functioning. Eur J Cardiothorac Surg 43: 95-103.

19. Aasvang EK, Gmaehle E, Hansen JB, Gmaehle B, Forman JL, et al. (2010) Predictive risk factors for persistent postherniotomy pain. Anesthesiology 112: 957-969.

20. Brandsborg B, Dueholm M, Nikolajsen L, Kehlet H, Jensen TS (2009) A prospective study of risk factors for pain persisting 4 months after hysterectomy. Clin J Pain 25: 263-268.

21. Nikolajsen L, Ilkjaer S, Krøner K, Christensen JH, Jensen TS (1997) The influence of preamputation pain on postamputation stump and phantom pain. Pain 72: 393-405.

22. Katz J, Asmundson GJ, McRae K, Halket E (2009) Emotional numbing and pain intensity predict the development of pain disability up to one year after lateral thoracotomy. Eur J Pain 13: 870-878.

23. Maguire MF, Latter JA, Mahajan R, Beggs FD, Duffy JP (2006) A study exploring the role of intercostal nerve damage in chronic pain after thoracic surgery. Eur J Cardiothorac Surg 29: 873-879.

24. Asch DA, Jedrziewski MK, Christakis NA (1997) Response rates to mail surveys published in medical journals. J Clin Epidemiol 50: 1129-1136.