



Open Access

## Prioritizing a Healthy Start: Nurturing Newborns through the Neonatal Period

a a a a a a a a a a .  
O a , / a / a a / a a a .  
a a b a a a a SIDS. Ma a  
a ab a a a a a a a .  
SIDS a a a a a a a a a .  
a a a a a a a a a a . E a a ab a  
a a a a a a a a a a .  
ba a a a b a a a a a . SIDS-  
a a a a . F a a a a a a a a a .  
a a a a a a a a a a . SIDS a a a a a a a a a a .  
a a a a a a a a a a . B  
a a a a a ab a a a a a -ba  
a a a a , / a a a a a a a a . SIDS a



2. Al Riyami N (2020) Respiratory distress syndrome in neonates delivered at term-gestation by elective cesarean section at tertiary care hospital in Oman. *Oman Med J* 17:133.
3. Tochie JN (2020) The epidemiology, risk factors, mortality rate, diagnosis, etiologies and treatment of neonatal respiratory distress: a scoping review.
4. Raha BK, MJ Alam, MAQ Bhuiyan (2019) নিম্ন গতিশীলতার জন্য পর্যবেক্ষণ করা হৃতি এবং পর্যবেক্ষণের ফল। *Bangladesh Medical Journal* 48: 21-27.
5. Raha BK, MJ Alam, MAQ Bhuiyan (2021) Spectrum of Respiratory Distress in Newborn: A Study From a Tertiary Care Military Hospital. *Bangladesh Coll Phys Surg* 39: 4-8.
6. পুরোজী এবং সুস্থ পর্যবেক্ষণের ফল। *A retrospective study on the risk of respiratory distress syndrome in singleton pregnancies with preterm premature rupture of membranes between 24+ 0 and 36+ 6 weeks, using regression analysis for various factors. Biomed Res Int* 15: 67-71.
7. Hermansen CL, KN Lorah (2007) Respiratory distress in the newborn. *Am Fam Physician* 76: 987-994.
8. Alfarwati TW (2019) Incidence, risk factors and outcome of respiratory distress syndrome in term infants at Academic Centre, Jeddah, Saudi Arabia. *Medical Archives*, 73: 183.
9. Sivanandan S, Agarwal R, Sethi A (2017) Respiratory distress in term neonates in low-resource settings. *Semin Fetal Neonatal Med* 22: 260-266.
10. Hubertus J, Plieninger S, Martinovic V, Heinrich M, Schuster T, et al. (2013) Children and adolescents with ureteropelvic junction obstruction: is an additional voiding cystourethrogram necessary? Results of a multicenter study. *Wor J Urol* 31: 683-687.