

Tgvtqrgtkvppgc"Uej ycppq o c<" Y j gp"GWU/ I wkfgf"HP C"ecp"Cxqkf"Uwtigt{

Claudio Zulli^{1*}, Nadia Alberghina¹, Giuseppe Grande¹, Mauro Manno¹, Luca Reggiani Bonetti², Flavia Pigò¹, Vincenzo Giorgio Mirante¹, Santi Mangiafico¹, Rita Conigliaro¹ and Carmelo Barbera¹

¹*Gastroenterology and Digestive Endoscopy Unit, NOCSAE Hospital, Baggiovara di Modena, Italy*

²*Department of Diagnostic Medicine and Public Health, University of Modena and Reggio Emilia-Section of Pathology, Modena, Italy*

***Corresponding author:** Claudio Zulli, Endoscopy Unit, NOCSAE Hospital, Baggiovara di Modena, Italy. Email: claudio.zulli@unimore.it




Figure 2 (C) EUS images of of the dis-homogeneous, hypoechoic, well-def ned, peripancreatic lesion. (D) EUS-guided FNA with 22G needle (no-styleset no-suction technique).



Figure 3 Biopsic fragments of tissue composed by spindle cells [Hematoxylin & Eosin stain; 4X]. G. Some fragments showed storiform pattern of growth [Hematoxylin & Eosin stain; 20X]. H: The spindle cells were positive for S100 [10X].