


## Risks and Outcomes of Extremity Fractures in Patients with Kntwith 湖レ昆虫

**Main Results:** Total participants from all these studies were 1,11,61,087  
participants in the past 10 years. The proportion of patients with mental health disorders was 10%.

**Conclusion:** Patients with fractures and underlying mental health disorders had longer hospital stay, poor functional outcomes and increased risk of chronic pain syndrome after discharge.



### Search terminologies

H (N L M ) (NLM) M  
(M H)  
F " OR M  
OR P D OR D I " AND M D  
D " OR " OR B D ". M H

### Inclusion criteria


A

74.3%


P I C D , N R , C  
M (ICD-9-CM) 12 6 13-18 ICD-  
10 14,15,19 4 20-23 . D  
D M M D , 5 2.5 , 21 . O 8036 , 2500  
(D M-5) 12,24 3 , 26 (31%) 1437 (17.7%) . I  
G M (AGECA ) 25 27 .

**Delayed discharge after fractures**


N 14-18,21,22,26,28 . H H  
(21.4% 2.2%, <0.001) 26 . P  
6  
M  
: 55%, : 41%, : 34%). P : 63%,  
( =0.001). P (31%)  
(24%) (33%) (25%). D (14%). H  
22 . O 822 , 14 (0.12%)  
I 28 (0.24%)



**Fracture risk in obese patients with mental health disorders**



**Pain perception & functional outcomes after fractures with mental health disorders**



**Discussion**

Discussion text describing the findings of the study, including the impact of mental health disorders on fracture risk and outcomes. The text discusses the association between mental health disorders and increased fracture risk, particularly in obese patients. It also addresses the impact of mental health disorders on pain perception and functional outcomes after fractures. The study highlights the need for a multidisciplinary approach to the management of these patients, involving both orthopedic and psychiatric professionals.

B  
L  
I  
E  
C  
J0  
9 0.2  
32  
44,45 . H  
K G P R D 17  
43  
42 . I

### Strengths and limitations of the selected articles

9-CM  
6  
4  
ICD-10  
H  
A  
26,28  
I

### Conclusions

P  
I

. I

. T



### Competing Interests

None.

### Contributorship

Both authors confirm that we had:

Substantial contributions to the conception and design of the work; the acquisition, analysis, and interpretation of data for the work; AND

Drafting the work and revising it critically for important intellectual content; AND

Final approval of the version to be published; AND

Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

### Funding Statement

This article received did not receive any grant from any funding agency, commercial or not-for-profit sectors.

### Ethical Approval Information

This article did not meet the criteria for obtaining an ethics committee or institutional review board approval as it does not reveal any identifiable personal, sensitive or confidential information of the participants of the included studies.

### Data Sharing Statement

Both the authors confirm that data file will be available to share on a reasonable request.

### References

1. Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, et al. (2007) Incidence and economic burden of osteoporosis-related fractures in the United States, 2005-2025. J Bone Miner Res 22: 465-475.
2. Reynolds CF, Cuijpers P, Patel V, Cohen A, Dias A, et al. (2012) Early

35. Lee SH, Hsu WT, Lai CC, Esmaily-Fard A, Tsai YW, et al. (2017) Use of antipsychotics increases the risk of fracture: a systematic review and meta-analysis. *Osteoporos Int* 28: 1167-1178.
36. Oderda LH, Young JR, Asche CV, Pepper GA (2012) Psychotropic-related hip fractures: meta-analysis of first-generation and second-generation antidepressant and antipsychotic drugs. *Ann Pharmacother* 46: 917-928.
37. Seppala LJ, Wermelink AMAT, de Vries M, Ploegmakers KJ, van de Glind EMM, et al. (2018) Fall-Risk-increasing drugs: A systematic review and meta-analysis: II. Psychotropics. *J Am Med Dir Assoc* 19: e11-e17.
38. Takkouche B, Montes-Martinez A, Gill SS, Etminan M (2007) Psychotropic medications and the risk of fracture: A meta-analysis. *Drug Saf* 30: 171-184.
39. Treves N, Perlman A, Kolenberg Geron L, Asaly A, Matok I (2018) Z-drugs and risk for falls and fractures in older adults-a systematic review and meta-analysis. *Age Ageing* 47: 201-208.
40. Pouwels S, Van Staa TP, Egberts AC, Leufkens HG, Cooper C, et al. (2009) Antipsychotic use and the risk of hip/femur fracture: A population-based case-control study. *Osteoporos Int* 20: 1499-1506.
41. Jalbert JJ, Eaton CB, Miller SC, Lapane KL (2010) Antipsychotic use and the risk of hip fracture among older adults afflicted with dementia. *J Am Med Dir Assoc* 11: 120-127.
42. Lewicki M, Paez H, Mandalunis PM (2006) Effect of lithium carbonate on subchondral bone in sexually mature Wistar rats. *Exp Toxicol Pathol* 58: 197-201.
43. Zamani A, Omrani GR, Nasab MM (2009) Lithium's effect on bone mineral density. *Bone* 44: 331-334.
44. Bolton JM, Metge C, Lix L, Prior H, Sareen J, et al. (2008) Fracture risk from psychotropic medications: A population-based analysis. *J Clin Psychopharmacol* 28: 384-91.
45. Vestergaard P, Rejnmark L, Mosekilde L (2005) Reduced relative risk of fractures among users of lithium. *Calcif Tissue Int* 77: 1-8.