Role of the TLR4 rs4986791 Polymorphism in the Development of Late-onset Alzheimer Disease and its Relationship with APOE*4

Romero R, Salazar Alcalá E, Cedeño N and Fernández-Mestre M*

Laboratorio de Fisiopatología, Centro de Medicina Experimental, Instituto Venezolano de Investigaciones Científcas, Caracas, Venezuela

Abstract

Objective: Based on that the association of the ε4 allele of the APOE gene with the development of late-onset AD is one of the strongest and that TLR4 has been involved in AD pathogenesis. The aim of the present work was to study the role of rs4986791 polymorphism of the TLR4 gene in the development of AD and correlate any such association with the presence of allele ε4 of the APOE gene.

Methods: We included 161 unrelated Venezuelan subjects classifed as either AD patients (n=61) or healthy individuals (n=100). Polymorphisms of TLR4 and APOE genes were identifed with PCR-SSP and PCR-RFLP, respectively.

Results: The rs4986791 polymorphism does not appear to be related to AD, although the presence of the CC genotype and the C allele apparently confers three times higher risk of developing AD. Finally, positive and negative associations among the combinations TLR4 /APOE genes and AD were observed.

Conclusion: The results suggest the absence of any association between rs4986791 polymorphism of TLR4 gene and susceptibility to AD and the association of the ε4 allele of the APOE gene with the development of this pathology was confrmed.

A speci c group of these receptors include the Toll-like receptors (TLRs). Toll like receptors belong to I transmembrane receptor family involved in the initial recognition of pathogens through speci c pathogen-associated molecular patterns (PAMP) receptors. e activation of TLRs triggers di erent signaling pathways, leading to the production of proin ammatory cytokines, reactive oxygen and nitrogen species [6]. Microglias express TLRs 1 to 9 and most of these receptors have been associated with its activation and neurotoxicity [2]. A characteristic in normal aging and in AD patients is an increase in innate immune receptors expression in the brain, such as TLRs. Particularly, high levels of CD14 (co-receptor for TLR4) in parenchymal microglia of the frontal and occipital cortex, hippocampus, and around the senile plaques have been observed in brain of AD patients [7]. Furthermore, monocytes derived from elderly individuals show defective TLR signaling and decreases in relative TLR1 and TLR4 expression [8]. TLR2 and TLR4 have been implicated in AD pathogenesis. A functional de ciency of either receptors results in an increased deposits of Aβ and reduced phagocytic clearance [5]. Speci cally, TLR4 has been associated with microglia activation, neurotoxicity and Aβ signaling. e investigations suggest that TLR4 play a dual role in the pathogenesis of AD. On the one hand, the TLRs are neuroprotective due to their contribution in the clearance of amyloid β. On the other hand, the in ammatory responses triggered by TLR4, through molecules associated with cell damage [5].

with those previously reported in the literature. A rst study showed that the 299Gly allele of the TLR4 gene may be associated with a decreased risk of late-onset AD in an Italian population sample, and this e ect was independent of the APOE e4 status [18]. Other studies showed a strong association between TLR4 polymorphisms and late-onset AD risk, especially among ApoE e4 non-carriers [19,20]. However, the mechanisms by which the polymorphisms in the TLR4 gene could a ect susceptibility to late-onset AD remain to be determined. In consequence future investigations should be focused on understanding the role of TLR4 genetic variants in neurodegenerative processes.

Allelic and genotypic frequencies of the APOE gene in healthy individuals and patients with Alzheimer disease

e distribution of the APOE allele and genotype frequencies in controls) and AD patients shows three common alleles (*APOE*2*, *APOE*3* and *APOE*4*). e *APOE*3* allele was the most frequent allele in both groups studied. However, the *APOE*4* allele frequency was signi cantly higher in patients with AD than in healthy individuals (37.7 % vs. 14.5% respectively, OR: 3.57, 95% CI: 2.0849-6.1093; *p*: 0.000001, *pc*: 0.000003) (Table 2). In addition, ve APOE genotypes were identi ed in patients and four in controls. A comparison of the frequencies of genotypes between AD and controls showed a signi cantly increased frequency of the ε3/ε4 (40.98% versus 21%, OR: 2.61; 95% CI: 1.2955-5.2677; *p*: 0.003, *pc*: 0.015) and ε4/ε4 (16.4% vs. 4%, OR: 4.7; 95% CI: 1.4058-15.7522; *p*: 0.008, *pc:* 0.04) genotypes in AD patients versus healthy individuals. Likewise, a signi cantly lower frequency of the ε2/ε3 (1.64% vs. 15%, OR: 0.094, 95% CI: 0.0121-0.7343, *p:* 0.003, *pc:* 0.012) and ε3/ε3 (39.34% vs. 60%, OR: 0.43, CI 95%: 0.2255-0.8292, *p:* 0.005, *pc:* 0.02) genotypes were recorded in patients versus controls (Table 2). Numerous studies have con rmed that the ε4 allele is the strongest genetic risk factor for sporadic AD. As compared to individuals with no ε4 alleles, the increased risk for AD is approximately is 2-3 fold in people with one ε4 allele and about 12-fold in those with two ε4 alleles [20,21]. ere are a variety of mechanisms by which apoE isoform may in uence risk for AD. Evidence indicates that the apoE isoforms di erentially a ect Aβ aggregation and clearance in the brain. However, other mechanisms likely to play a role in the ability of apoE to in uence CNS (Central Nervous System) function

Genotypes APOE	AD (n=61)	Control (n=100)	OR 95% CI	p
ε2/ε3	1.64 (1)	15 (15)	0.094 (0.0121-0.7343)	0.003
ε2/ε4	1.64 (1)	0	4.98 (0.1998-124.2621)	ns
ε3/ε3	39.34 (24)	60 (60)	0.43 (0.2255-0.8292)	0.005
ε3/ε4	40.98 (25)	21 (21)	2.61 (1.2955-5.2677)	0.003
ε4/ε4	16.4 (10)	4 (4)	4.7 (1.4058-15.7522)	0.008
Alleles				

as well as AD. ese mechanisms include e ects on synaptic plasticity, cell signaling, lipid transport and metabolism, and neuroin ammation [22].

Study of the combined e ect of polymorphism in TLR4 and APOE genes

ere of the 12 possible combinations of *TLR4/APOE*, seven were observed in both groups. e combinations CC/ε*3*ε*4* (OR: 2.6; 95% CI: 1.2757-5.2766; *p:* 0.003; *p*c: 0.021) and *CC/*ε*4*ε*4* (OR: 4.7; 95% CI: 1.4058-15.7522; *p*: 0.008, *p*c: 0.05) were signi cantly more common in AD patients than in controls (associations positive). We must point out that some genotype combinations were more common in the control group: *CC/*ε*2*ε*3* (OR: 0.11, 95% CI: 0.0142-0.8753, *p*: 0.009, *p*c: not signi cant) and *CC/*ε*3*

Citation: Romero R, Salazar Alcalá E, Cedeño N, Fernández-Mestre M (2018) Role of the TLR4 rs4986791 Polymorphism in the Development of Late-onset Alzheimer Disease and its Relationship with APOE*4. J Alzheimers Dis Parkinsonism 8: 436. doi: 10.4172/2161-0460.1000436