Keywords: Senegal; Seroprevalence; Influenza A virus subtype H1N1; Risk factors

Introduction

In June 2009, the WHO declared the influenza A (H1N1) pdm09 virus outbreak as the first pandemic of the 21st century. It reached more than 200 countries and was responsible for over 18,000 deaths worldwide [1]. In Africa, the A (H1N1) pdm09 virus was detected for the first time in Cape Verde in June 2009 [2]. In May 2010, the WHO announced more than 18,000 confirmed cases identified by local monitoring networks [3].

In Senegal, A (H1N1) pdm09 virus was detected by the influenza sentinel surveillance network entrusted to the Pasteur Institute of Dakar (IPD) [4] for the first time in January 2010 with a single epidemic peak in February 2010 [2]. Key collected blood samples taken from symptomatic individuals who had consulted a sentinel health center spontaneously. Key identified 345 cases of A (H1N1) pdm09/1328 samples indicating epidemic [illegible]

Variables	n	%
Gender		
Men	1038	39%
Women	1622	61%
Age		
[2-6]	220	8.3%

threshold, the rate of positive A (H1N1) pdm9 decreases slowly, from 74.5% to 10.0% (Figure 1).

A (H1N1) pdm09 seroprevalence di ered signi cantly between neighborhoods (p<0.001). e lowest prevalence was in Baghdad neighborhood, 16.8% (12.8-20.7) and the highest was in Soce Sow neighborhood, 71.0% (66. 7-75.3) (Figure 2).

Individual socio-demographic variables and health data reported

ere was no signi cant association between either sex or ethnicity and A (H1N1) pdm09 serology. Age class was globally signi cantly associated with A (H1N1) pdm09 positivity (p global=0.03). Only the 2-5 years age group had a signi cantly higher positivity than the reference class (25-44 years) (OR=1.83 (1.25-2.69)). e link between A (H1N1) pdm09 serology and variable “main contact type” was close to signi cance (p=0.056). Among people who reported a history of u over the past three years, only 39.9% were positive for A (H1N1) pdm09.

Variables related to household composition

ere was no signi cant association between A (H1N1) pdm09 serology and (i) the number of people living in the household (p=0.27), (ii) the number of children in the household (p=0.46), (iii) the number of school attending children living in the household (p=0.91).

Protective measures against A (H1N1) pdm09 u

e variable “how many times did you wash your hands with soap yesterday” was coded into three classes (0 times, 1-2 times or 3 times per day) and was signi cantly associated with a reduction in A (H1N1) pdm09 infection (p global=0.03).

Regarding household variables, “living in household having adopted hand washing” was not associated with decrease of A (H1N1) pdm09 infection (OR=0.72 (0.48-1.07); p global=0.11). However, “living in a household having adopted anti-cough prevention method” was signi cantly associated with a decrease of A (H1N1) pdm09 infection (OR=0.44 (0.24-0.84), p=0.01).

Multivariate analysis

We included in the multivariate model variables with p 0.20 in the bivariate analysis: “type of main contact”, “hand washing the day before”, “household having adopted home hand washing “,” household having adopted anti-cough prevention method”, “presence of a smoker at home” (Table 2). We chose not to include age in the model because of its

14. Nicolas S (2013) Contribution to the sero-epidemiological study of the fu. Thesis: specialty communicable diseases and tropical parhologies. University of Marseille 138.

15. Demographic and Health Survey of Multiple Indicators in Senegal (EDS-MICS) 2010-2011. National Agency of Statistics and Demography (ANSD), Senegal 2012.

16. Delabre RM, Lapidus N, Salez N, Mansiaux Y, De Lamballerie X, et al. (2015) Risk factors of pandemic infuenza A/H1N1 in a prospective household cohort in the general population: results from the CoPanFlu-France cohort. Infuenza Respir Virus 9: 43-50.

17. Niang MN, Dosseh A, Ndiaye K, Sagna M, Gregory V, et al. (2012) Sentinel Surveillance for Infuenza in Senegal, 1996-2009. J Infect Dis 1: 129-135.

18. Miller E, Hoschler K, Hardelid P, Stanford E, Andrews N, et al. (2009) Incidence of 2009 pandemic infuenza A H1N1 infection in England: a cross-sectional serological study. Lancet 375: 1100-1108.

19. Broor S, Krishnan A, Roy DS, Dhakad S, Kaushik S, et al. (2012) Dynamic

Pa /Ac 7/2 /Ta /Ar /Ty /Ex /Le /Ma /Ob /Ty /Ex /Le /Ma /Ob /Ty /Ex /Le /Ma /Ob