

# Sexual and Reproductive Health Utilization and Satisfaction among Adolescents in Northern Nigeria: Impact of REACH Project Educational Intervention

Yakubu Lawali<sup>1\*</sup> and Rahinatu Adamu Hussaini<sup>2</sup>

<sup>1</sup>Department of Nursing Sciences, Usmanu DanFodiyo University Sokoto, Sokoto, Nigeria

<sup>2</sup>Save the Children International, Abuja, Nigeria

## Abstract

**Objective:** The aim of the study was to evaluate the impacts of Reaching and Empowering Adolescents to make informed Choices for their Health (REACH) project intervention in utilization and satisfaction of Adolescents' Sexual and Reproductive Health (ASRH).

**Methodology:** This was a mixed methods study which collected both quantitative data (via a structured household survey and a health facility survey) and qualitative data (via Focus Group Discussions (FGDs) and Key Informant Interviews (KIIs)). The primary study population was male and female adolescents aged between 10 and 19 years within the three states (Gombe, Katsina and Zamfara).

**Results:** The finding of this project shows that, there is an increase in utilization of SRH services by adolescents as compared to the base line. Similarly, the result shows that married girls were more likely than unmarried girls to report using SRH services: 86% and 44% respectively ( $\chi^2$   $p < 0.001$ ). The same was true of boys, but the difference between married and unmarried boys was much smaller: 48% and 40% respectively. Similarly, there was an increase in the satisfaction with ASRH services as compared to base line.

**Conclusion:** REACH project educational intervention is effective in increasing utilization of Adolescents Sexual and Reproductive Health (ASRH) services.

**Keywords:** Adolescents • Adolescents' sexual and reproductive health • Structured household survey • Health facility survey

**Abbreviations:** REACH: Reaching and Empowering Adolescents to make informed Choices for their Health; LGA's: Local Government Authorities; ASRH: Adolescents Sexual and Reproductive Health; EL: End Line; BL: Base Line.

## Introduction

The World Health Organization (WHO) defines "adolescents" as individuals in 10–19 years old and "youth" as 15–24 years old [1]. Together, adolescents and youth are referred to as young people, encompassing the ages of 10–24 years. Studies have established what can increase access and utilization for SRH among adolescents. However, a lack of scientifically sound data on the effectiveness of services that target young people in sub-Saharan Africa, especially in comparison to the magnitude of Adolescents Sexual and Reproductive Health (ASRH) challenges in the region [2]. There is a gap in adolescents' access to Sexual and Reproductive Health (SRH) services and information, which has not been fully addressed. In addition, adolescents have now been included in the World Health Organization's Global strategy for women's, children's and adolescents' health (2016–2030), and this indicate the unique health challenges facing young people [3]. This is one of the reason for the implementation of REACH project by the save the children International in Nigeria.

The Reaching and Empowering Adolescents to make informed Choices for their Health (REACH) project was funded by Global Affairs Canada and

implemented by Save the Children (SC) between April 2018 and August 2021. Over its three years of implementation, it aimed to improve the Sexual and Reproductive Health (SRH) of adolescent boys and girls aged 10–19 within three Nigerian states: Gombe, Zamfara and Katsina.

REACH aimed to increase accessibility to high-quality and gender-responsive Adolescent Sexual and Reproductive Health (ASRH) services

Children decided that the MTR should form the basis for the EL evaluation, and that the EL data collection should consist of some targeted qualitative work to assess:

- The elements of the project which were implemented in its later stages.
- The extent to which the MTR's main recommendations had been implemented during the final months of the project. This report contains data from both the MTR and the additional EL qualitative work, which taken together form the full EL evaluation.

---

## Methodology

This was a mixed methods study which collected both quantitative data (via a structured household survey and a health facility survey) and qualitative data (via Focus Group Discussions (FGDs) and Key Informant Interviews (KIIs)). The primary study population was male and female adolescents aged between 10 and 19 years within the three states (Gombe, Katsina and Zamfara). The study also covered key influencers: parents of adolescent girls, adult husbands of adolescent girls, health facility staff, State and Local Government Area (LGA) administrators, Patent and Proprietary Medicine Vendors (PPMVs), traditional and religious leaders, and REACH facilitators.

### Sample size and sampling strategy

**Household survey:** The survey used a multi-stage cluster sample with 4 stages: state, LGA, village/community and individual. All three REACH states were selected: Gombe, Katsina and Zamfara. Within each state, two LGAs were selected, making a total of six LGAs.

Villages/communities were sampled at random from the list of catchment communities located within 5 km of the programme's target health facilities. Initially two villages/communities were selected in each of the 6 LGAs, but there were very few married 10-14 year-olds in these villages/communities, so two additional ones were selected in the two LGAs in Zamfara state. Thus, 14 communities were selected, with the number of communities selected per LGA approximately proportional to the size of the target population. Two of the 14 communities were also included in the BL sample (Table 1).

The aim was to conduct approximately 400 interviews per state (1,200 in totals); including adolescents aged 10-19, parents of adolescent girls and adult husbands of adolescent girls. Households were selected using a random walk method, with enumerators stopping at every second or third household (depending on the size of the village/community). At each sampled household, enumerators asked if any eligible respondents were resident and available at the time of the visit. If so, the interview was conducted immediately. If an eligible household was identified but the eligible respondent was not available when the enumerator visited, the enumerator returned at a different time of day to conduct the interview. If the eligible respondent was still not available at the return visit, the household was replaced by another, using the same sampling method.

**Adolescents:** To be eligible for interview, an adolescent had to: be aged 10-19 (inclusive) on the day of the interview, be resident in one of the sampled communities and have benefited from the REACH programme.

Enumerators were instructed to ensure that, within each age and sex group (boys 10-14, girls 10-14, boys 15-19 and girls 15-19) the sample included both married and unmarried adolescents. In the event, even with the addition of two new villages/communities (see above), the enumerators were not able to find many married 10-14 year-olds who met the inclusion criteria. They therefore supplemented the sample with additional married 15-19 year-olds, to ensure that the sample contained sufficient married adolescents for separate analysis. This deliberate over-sampling of married adolescents is almost certainly the

---

---

## Discussion

The finding of this project shows that, there is an increase in utilization of SRH services by adolescents as compared to base line. Specifically, there was increased in utilization of family planning services with 32.2%. Similarly, the result shows that married girls were more likely than unmarried girls to report using SRH services: 86% and 44% respectively ( $p < 0.001$ ). The same was true of boys, but the difference between married and unmarried boys was much smaller: 48% and 40% respectively. The greater use among married girls than among married boys reflects the attitudes described earlier about husbands tending to make the FP decisions and wives tending to put them into practice. These findings are similar to that of Banke-Thomas and Ameh [5,6].

The BL assessment found that 58% of married girls and 47% of married boys said they had received SRH services in a health facility. The equivalent EL figures of 86% and 44% respectively indicate a major increase in married girls' uptake of adolescent SRH services, but no significant change among married boys [7-10].

---

## Conclusion

Among the 423 adolescents in the household survey who said they had used SRH services, nearly all (96%) expressed satisfaction with these services. On this note, the adolescents reported 28% increase in the utilization of these services. The BL assessment found that 66% of married girls and 70% of married boys were satisfied with the SRH services in health facilities.