



• MRI analysis; Brain abnormalities; Language and behavioral disorders; Neurolinguistics; Bangladesh

Habitant		
Rural	42	56
Urban	33	44
Parents education		
Illiterate	43	54
Primary (class-V)	21	22
Secondary (class-VIII)	7	14
SSC	4	10

School Certificate.

Ferdous F, Monte-Serrat DM, Nabi S (2024) Significance of Neurolinguistic Approach to Brain Abnormalities Perceived by Magnetic Resonance Imaging (MRI) among Children with Language and Behavioral Disorders: A Cross Sectional Descriptive Observational Study. *J Alzheimers Dis Parkinsonism* 14: 592

1. Chin EM, Jayakumar S, Ramos E, Gerner G, Soares BP, et al. (2019) Preschool language outcomes following perinatal hypoxic-ischemic encephalopathy in the age of therapeutic hypothermia. *Dev Neurosci* 40(5-6):627-637.
2. Martinez-Biarge M, Diez-Sebastian J, Rutherford MA, Cowan FM (2010) Outcomes after central grey matter injury in term perinatal hypoxic-ischaemic encephalopathy. *Early Hum Dev* 86(11):675-682.
- 3.
12. Max JE, Keatley E, Wilde EA, Bigler ED, Schachar RJ, Saunders AE, et al. (2012) Depression in children and adolescents in the first 6 months after traumatic brain injury. *Int J Dev Neurosci* 30(3):239-245.
13. Shapiro KA, Kim H, Mandelli ML, Rogers EE, Gano D, et al. (2017) Early changes in brain structure correlate with language outcomes in children with neonatal encephalopathy. *Neuroimage Clin* 15:572-580.
14. Steinman KJ, Gorno-Tempini ML, Glidden DV, Kramer JH, Miller SP, et al. (2009) Neonatal watershed brain injury on magnetic resonance imaging correlates with verbal IQ at 4 years. *Pediatrics* 123(3):1025-1030.
15. Reynolds JN, Wickens JR (2002) Dopamine-dependent plasticity of corticostriatal synapses. *Neural Netw* 15(4-6):507-521.
16. Martinez-Biarge M, Bregant T, Wusthof CJ, Chew AT, Diez-Sebastian J, et al. (2012) White matter and cortical injury in hypoxic-ischemic encephalopathy: antecedent factors and 2-year outcome. *J Pediatr* 161(5):799-807.
17. Van Agtmaal MJ, Houben AJ, De Wit V, Henry RM, Schaper NC, et al. (2018) Prediabetes is associated with structural brain abnormalities: the Maastricht Study. *Diabetes care* 41(12):2535-2543.
18. Ferdous F, Monte-Serrat DM, Chisty MM, Alam MF (2017) A fourth-generation methodological practice on a study of speech dysfunction of children with cerebral palsy. *Int J Percent Publ Health* 2(1):11-23.
19. Hinde K, Mason J, Kannis-Dymand L, Milleer P, Sultana R (2023) Bangladeshi Health Practitioner Knowledge, Confidence in Diagnosis, and Treatment of Mental Health Disorders in People with Intellectual Disabilities. *J Ment Health Res Intellect Disabil* 16(3):205-225.
20. Martinez C, Carneiro L, Vernier L, Cesa C, Guardiola A, et al. (2014) Language in children with neonatal hypoxic-ischemic encephalopathy. *Int Arch Otorhinolaryngol* 255-259.
21. Liégeois F, Mayes A, Morgan A (2014) Neural correlates of developmental speech and language disorders: evidence from neuroimaging. *Curr Dev Disord Rep* 1(3):215-227.
22. Bax M, Tydeman C, Flodmark O (2006) Clinical and MRI correlates of cerebral palsy: the European Cerebral Palsy Study. *JAMA* 296(13):1602-1608.