

/s

the first two digits of the number of the sample.

■ ■ ■

After the samples were taken, the soil was washed with water and the residue was dried at 100°C for 24 hours. Then the residue was weighed and the percentage of sand was calculated. The results are shown in Table I.

The results show that the percentage of sand in the samples taken from the surface layer ( $0-10$  cm) is higher than that in the samples taken from the deeper layers. This indicates that the sand content in the surface layer is higher than that in the deeper layers.

The results also show that the percentage of sand in the samples taken from the surface layer ( $0-10$  cm) is higher than that in the samples taken from the deeper layers. This indicates that the sand content in the surface layer is higher than that in the deeper layers.

The results also show that the percentage of sand in the samples taken from the surface layer ( $0-10$  cm) is higher than that in the samples taken from the deeper layers. This indicates that the sand content in the surface layer is higher than that in the deeper layers.

The results also show that the percentage of sand in the samples taken from the surface layer ( $0-10$  cm) is higher than that in the samples taken from the deeper layers. This indicates that the sand content in the surface layer is higher than that in the deeper layers.

After the samples were taken, the soil was washed with water and the residue was dried at 100°C for 24 hours. Then the residue was weighed and the percentage of sand was calculated. The results are shown in Table I.

The results show that the percentage of sand in the samples taken from the surface layer ( $0-10$  cm) is higher than that in the samples taken from the deeper layers. This indicates that the sand content in the surface layer is higher than that in the deeper layers.

The results also show that the percentage of sand in the samples taken from the surface layer ( $0-10$  cm) is higher than that in the samples taken from the deeper layers. This indicates that the sand content in the surface layer is higher than that in the deeper layers.


The results also show that the percentage of sand in the samples taken from the surface layer ( $0-10$  cm) is higher than that in the samples taken from the deeper layers. This indicates that the sand content in the surface layer is higher than that in the deeper layers.

The results also show that the percentage of sand in the samples taken from the surface layer ( $0-10$  cm) is higher than that in the samples taken from the deeper layers. This indicates that the sand content in the surface layer is higher than that in the deeper layers.

The results also show that the percentage of sand in the samples taken from the surface layer ( $0-10$  cm) is higher than that in the samples taken from the deeper layers. This indicates that the sand content in the surface layer is higher than that in the deeper layers.

1 /s /s 1 /s







31. Tran LSP, Nakashima K, Sakuma Y, Simpson SD, Fujita Y, et al. (2004) Isolation and functional analysis of *Arabidopsis* stress-inducible NAC transcription factors that bind to a drought responsive cis-element in the early