treated rabbit erythrocytes. e hemagglutination titre was obtained to be 5 × 10 hemagglutination units per mg of protein. Simple sugars like glucose, mannose, galactose, rhamnose, xylose, fucose, ra nose, trehalose, lactose, glucosamine, mannosamine, galactosamine, methyl- -glucose, methyl- -mannose, methyl- -galactose, N-acetyl-mannosamine, N-acetyl-galactosamine, N-acetyl-D-glucosamine, D-mannitol, D-glucuronic acid, D-polygalacturonic acid and D-glucose pentaacetate failed to inhibit the hemagglutinating activity of the lectin. Hemagglutination by rCAL was inhibited by glycoproteins viz. fetuin, asialofetuin, and its asialotriantennary glycans. e minimum inhibitory concentration for fetuin and its asialo glycan was found to be 32 µg. Lectins recognizing only complex glycans have been reported earlier. For example, the lectin from wild cobra, Naja naja avum [22] shows inhibition only for asialofetuin with an inhibitory concentration of 250 µg/ml.

Ligand binding with Asialo triantennary glycan from Fetuin

Fluorescence titration of the rCAL with N-linked asialo triantennary glycan prepared from fetuin structure shown (Figure

residues per chain)), LYS-335 (B chain - 107), ALA-337 (B chain - 106), ASP-338 (B chain - 109) PRO-341 (B chain - 110) and GLU-344 (B chain - 113) (Figure 3C). e most favourable docked pose of hemin had a binding energy of -31.35 kJ^{-1}. For spermine and thiamine also, the binding sites on the model were present only on one of the two monomers and located close to each other (Figure 4A). Spermine interacted with amino acids ASN-8 (2.1 Å), ASP-66 (2.1 Å) and ASP-122 (1.9 Å) of rCAL by formation of hydrogen bonds (bond lengths given in brackets) (Figure 4B). iamine formed hydrogen bonds with ASP-66 (2.2 Å), TYR-80 (1.9 Å) and LYS-137 (2.2 Å) of rCAL (Figure 4C). Binding energies of -15.88 kJ^{-1} and -22.15 kJ^{-1} were obtained for spermine and thiamine, respectively.

Interestingly, both hemin and spermine were found to interact with ASN-8 of the protein. Gaur et al. [10] have shown that hemin and spermine bind to the LS-24 albumin in a mutually exclusive manner. Spermine is bound to the dimeric LS-24 albumin, but hemin binding results in dissociation of the subunits accompanied by loss in the ability to bind spermine. is binding property might be true for rCAL also, wherein a common residue is involved in the binding of both hemin and spermine.



ASP-122

In the crystal structure of LS-24, binding of spermine involves the interaction of GLU-80, ASN-81, and SER-118 [10]. In the rCAL model, spermine binding involves ASN and S04e band ais is.-30(a)9(e)s-3SiMl e r aASN albuaN apermine. Tfbr(n)4(d)1m(ua)-6pBad a 18(n)4(dx