human experience [20-22]. Through story-telling meaning is constructed from the experience [21,23].

In narrative inquiry, the interview should be approached with the intent of hearing stories. Narrative has a commitment to the value of giving voice to human feelings and experience [24], through the use of "imaginative reconstruction" or "empathy" [25].

Grinyer [26] believes narrative is an appropriate method for sensitive research as it allows family members to determine the content of the information they share which she sees as strength rather than a weakness.

Narrative inquiry also includes the concept of a shared reality [20]. Bruner [20] has said that "life stories must mesh tellers and listeners must share some 'deep structures' about the nature of a life." If this does not occur, both the storyteller and the listener will fail to grasp what the other is telling or hearing.

Narrative Inquiry therefore is a suitable methodology for this research study. Within its epistemology, understanding is the development of knowledge through the telling of stories. These stories create meaning and a testament of the experience of suffering, giving voice to what might not be told.

The methodology suggests my own experience, rather than hindering the research, creates the context for me as the researcher to grasp what the family member is saying and provide a richer understanding of what is being told.

Ethics approval was obtained through the Southern Cross University Human Research Ethics Committee (Approval number ECN-05-146).

The family members in this study were a purposeful sample of 26 who had experienced the death of an AYA (aged 13 to 23) to cancer. Table 1 provides a representation of the family members relationship to the AYA, the interview type and time since death. The interviews were open-ended. All interviews were recorded on an iPod and professionally transcribed.

For the purpose of this study, family members preferred not to use pseudonyms and requested that their family member's real names be used. Additional ethical approval was obtained from Southern Cross University's Human Research Ethics Committee for this to be possible (approval number ECN-08-029).

The study utilised narrative analytic methods to create a configured plot of core stories and a configured meta-narrative from which commonalities and differences within the core stories were identified.

Guidance was obtained from the following narrative theorists: Mishler [27], Riessman [28,29], Polkinghorne [30], Lieblich, Tulval & Zilber [31], Emden [21,32], and Frank [33], to analyse these narratives.

In addition Taylor's [34] computer-assisted method and the qualitative research software program NVivo8 provided a framework to configure and organise the stories into a meta-narrative.

Figure 1 links the steps of the narrative analysis for this study with specific processes and relevant authors.



Figure 1: Steps to Narrative Analysis.

The narrative patterns within the stories identified transitional or turning points in the family members' stories.

T is article provides insights into the experience of family members through themes identi fed

Recalling the moment	17	46
Waiting and watching Being present Missing the moment Sensing relief		
Extending the living connection	11	30
Staying with the body Caring for the body Viewing the body		

Table 1: T e Dying Landscape.

T e results are a summary of the family members' stories within the themes. Direct quotes from the participants stories are represented by quotation marks and italics.

T e death and dying stage for the AYAs and family members in this study commenced when the AYA exhibited signs and symptoms that alerted the AYA and the family members that things were not getting better or the cancer had returned. Denise was shocked when Brenton James's melanoma reoccurred in the liver as they had been told he was f ne. She said *"You know we got through this shocking period of him losing his eye and we'd been given the all clear. We'd got through this, we'd escaped."* Lauren noticed that Grant was not getting any better which was frustrating *"then in the April he started coughing and we called the Oncologist and they said it is probably just a virus … but we will get an x-ray just in case and that is when we found the tumours in his lungs, So that was pretty devastating."* For most, tests and investigations were quick to con f rm the return of the cancer. For Tamina who had transitioned from the paediatric to the adult system there was evidence of lack of communication within these two systems, as was seen in one story that resulted in a delay in diagnosis. Pam said *"we've been coming back and forth, or Tamina had been for a few months with this cough … She was told she had a lung infection … and nobody cottoned on to actually send her for a chest x-ray."* When a Chest X-ray was f nally ordered they found she had relapsed.

T e

T e way that the family members were told of the impending death was not always satisfactory. Denise spoke of the frustration she felt when a young doctor tried to tell her Brenton James was dying: *"the intern doctors … a lot of them are still f nding their feet with how to deal with these things … especially the one that was trying to tell me that my son was critical and that he would die, he rambled on and on and on and on, and I wanted to shake him. To say … just say it … [I] know what you're trying to say, but I didn't."*

T ere was no evidence of a plan as to when, where, and how the family members were told, added to this there was little support for the AYA and the family member a f er the pronouncement. T is is seen in Jenny's story, she said the doctor rang her at work and said*: "I'm sorry, but Brenton is going to die, we can't do anything about it. He has got metastasis everywhere".* Jenny then had the responsibility of telling her



son, Brenton Duncan that he was going to die. For Pam, Tamina was told she was going to die with no support from family. Pam said *"she (Tamina) was in there for a while [the doctors surgery] and he told her she was going to die … and they knew I was out in the waiting room."* Understandably the response for most was shock and devastation. Lauren recounted *"the fact that that's it, there is nothing more we can do for you was horrible."* For Fulvia the pronouncement of impending death did not occur until the AYA was near death. For example Fulvia recalled a doctor describing the dying process. She says doctor said: *"Jadye's condition has deteriorated, Jadye's now unconscious and soon she's going to start cheyne stoking, she's going to start breathing di erent and so on and I don't want you all to be distressed, it's just the normal way of dying, there's no pain and she's not in any distress."*

T e family members noticed that with the pronouncement of impending death the focus of care changed both in how and where the care was delivered. Lauren spoke of how this quality of care changed: *"whereas in the f rst time it was, well it doesn't matter how sick he gets because in the end … he'll be alive. … and so that really changed … it de f nitely didn't r e r @ 'sus'lu'¢im' at"std] te! e m#h*

going to go through this, stay at home if you can get treatment at home rather than be in a hospital."

Staying home for the last moments was not always a choice that the

tube in, say his goodbyes and then we would take the tube out and he

needs during and after the death and to see the person as not just a body. Whilst there was a sense of relief at the moment of death, this moment augurs the transition to another stage incorporating the loved one into their life with memories and rituals.

Barling [8] sought to understand the experience of the AYA cancer journey from the perspective of family members. This article presents selected findings from the Barling [8] study that explained family members' experience of during the destinations stage of the cancer journey.

The summary of the meta-narrative provides acknowledgment that the destination stage is difficult for the family members and the AYA. The stories reveal a lack of support and being overwhelmed with the burden of care as the family members maintained their vigil. Despite this the family members found it important to talk about the death and how they continued their living connection with their loved one after death. In addition, the absence of EOL discussions within the majority of the meta–narratives suggests that AYAs and family members are often not provided with the opportunity to consider EOL choices during the destination stage.

This article has therefore highlighted the need to continue research into the needs of AYAs and their families during the destination stage to further increase understanding and assist in the implementation of the National and OnTrack Framework recommendations, with particular emphasis on specific issues associated with the destination stage of the cancer journey.

35. Armstrong-Coster A (2004) Living and dying with cancer. Cambridge: Cambridge University Press.
36. De Jong J, Clarke L (2009) What is a good death? Stories from palliative care. J Palliat Care 25: 61-64.
37. Bell C (2010) End of life experiences in adolescents dying with cancer. Support Care Cancer 18: 827-835.
38. Foreman L (2006) Factors predictive of preferred place of death in the general population of South Australia. Palliat Med 20: 447-453.
39. Grinyer A, Tomas C (2004) The importance of place of death in young adults with terminal cancer. Mortality 9: 114-131.
40. Hinds P, Pritchard M, Harper J (2004) End of life research as a priority for pediatric oncology. J Pediatr Oncol Nurs 21: 175-179.
41. Stillion J, Papadatou D (2002) Sufer the children: An examination of psychosocial issues in children and adolescents with terminal illness. American Behavioural Scientist 46: 299-315.
42. Williams A (1998) Therapeutic landscapes in holistic medicine. Soc Sci Med 46: 1193-1203.
43. Brown P, Davies B, Martens N (1991) Families in supportive care - Part 1: Palliative care at home a viable care setting. J Palliat Care 6: 21-27.
44. Clayton J (2007) Clinical practice guidelines for communicating prognosis and end -of life issues with adults in the advanced stages of a life-limiting illness, and their caregivers. Med J Aust 186: 77-108.
45. McGrath P (2002) End of life care for hematological malignancies : Technological imperative and palliative care. J Palliat Care 18: 39-47.
46. McGrath P (2001) Dying in the curative system: The haematology/ oncology dilemma. Part 1. Aust J Holist Nurs 8: 22-30.
47. McGrath P (2002) Dying in the curative system: The haematology/ oncology dilemma. Part 2. Aust J Holist Nurs 9: 14-21.
48. Wittenberg-Lyles E, Goldsmith J, Ragan S (2011) The shif to early palliative care: : a typology of illness journeys and the role of nursing. Clin J Oncol Nurs 15: 304-310.
49. Wolfe J (2000) Symptoms and sufering at the end of life in children with cancer. N Engl J Med 342: 326-333.
50. (2009) CanTeen and Cancer Australia National Service Delivery Framework for Adolescents and Young Adults with Cancer.