

Substance Use Disorder: Signs and Symptoms

Michael Stevens *

Department of Psychiatry, University of Florida, USA

Corresponding author:

Current problems and costs, economically, medical, legal and personal, local pain medications and chronic pain treatment are never ending.

6XEVWDQFH XVH GLVRUGHU LV GUXJ -XVH GHVSLWH WKH VLGH HuHFWV 7KLV FRQ WLRQ LQFOXGHV GUXJ UHVLVWDQFH KLJK FUDYLQJ IRU GUXJV DQG GLvFXOW\ L

reducing drug use. Withdrawal symptoms may include insomnia, nausea, muscle aches, and diarrhea. Diacetylmorphine is used as a mind-

DOWHULQJ GUXJ ,W DOVR LQFOXGHV VLGH HuHFWV VXFK DV VKRUWQHVV RI EUH

drowsiness, and addiction. Its therapeutic objectives include pain medications such as trauma, postoperative pain or heart disease. Physical symptoms include stretching, yawning, sneezing, tremors, high pain,

high blood pressure, fever, light sensitivity, restlessness and vomiting. Drug overdoses, both lethal and non-lethal are common. Most homicides happen at home, with a few seemingly intentional. Recent studies

show that disturbed breathing during non-REM sleep increases with RSLRLG GRVDJH 7KH SRZHUIXO HuHFW RI WKH GUXJ RQ UHVSLUDWRU\ GLVWUHV

between both animals and humans is well documented. Substance use disorder impairs perception and emotional distress, and contributes to poor judgment. Forced use due to addiction leads to loss of control and

GDQJHURXV XVH RI WKH GUXJ DQG VLGH HuHFWV %XSUHQRUSKLQH XVHG WR WU SDLQ DQG DGGLFWLRQ 6LGH HuHFWV- RI EXSUHQRUSKLQH LQFOXGH QHXUDO LQKI

WLRQ SRVWXUDO K\SRWHQVLRQ ZHLJKW ORVV GU\ PRXWK DQG HuHFWV RQ W FHQWUDO QHUVRXV V\VWHP PHPRU\ ORVV DQG GL]]LQHVV %HQ]RGLD]HSLQH

are commonly prescribed to treat opioid-related anxiety and depression.

When a medical professional writes these instructions, patients and

SK\VLFLDQV DOLNH RIWHQ IDLO WR P-DNH VLJQL¿FDQW LQWHUDFWLRQV DQG LQW WLRQV EHWZHHQ WKHVH GUXJV DQG WKH RSSRVLWH HuHFWV DYDLODEOH \$OWK

there is evidence that severe pain from opioid medications in the short

WHUP WKHUH LV QR VLJQL¿FDQW HYLGHQFH RI LPSURYHG SK\VLFDL DFWLYLW\

maintenance of pain relief for long-term remission. Methods of loss of DQDOJHVLF HvFDF\ LQFOXGH WKH GHYHORSHPHQW RI SKDUPDFRORJLF WROHUDQF

and opioid-induced hyperalgesia, addiction and dependence. Pain management with opioid medications is a commonly discussed and related