

Sugar Partitioning and Sink-Source Modification in Plants

Mohammad Reza Siahpoosh*

Department of Agronomy and Plant Breeding, Shahid Chamran University, Ahwaz, Iran

*Corresponding author: Mohammad Reza Siahpoosh, Department of Agronomy and Plant Breeding, College of Agriculture, Shahid Chamran University, Ahwaz, Iran, Tel: +98 61 3330010-20; Email: siahpoosh@scu.ac.ir

Rec date: Sep30, 2014, Acc date: Oct 10, 2014, Pub date: Oct 15, 2014

Copyright: © 2014 Siahpoosh. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Editorial

Solar energy is converted to chemical energy and stored as assimilates through a phenomenon called photosynthesis. Plant leaves function as the principle site of resource acquisition by utilizing the free energy captured via photosynthesis for the reductive assimilation of oxidized forms of carbon into carbohydrates. Photosynthetic carbon fixation provides vital energy for metabolism and precursors for all other biosynthetic pathways in the plant. Most of these precursors are required for biosynthesis of amino acids that form the building blocks for many compounds in plants. The regulation of assimilate partitioning in leaves is considered as allocation of carbon between sucrose and starch synthesis, storage, and export, and carbohydrate metabolism [1,2]. Sucrose is the most important metabolite in this system of resource allocation because it is generally the major end product of photosynthetic carbon metabolism and, in most plants it is the predominant form of carbon transported to the heterotrophic tissues [3-5]. Sucrose allocation between tissues is a fundamental process in all multicellular organisms. Indeed, as much as 80% of the carbon acquired in photosynthesis is transported in the plant's vascular system to import-dependent organs [6].

Moreover, in many plants, energy-dependent sucrose accumulation in the phloem generates the high hydrostatic pressure that drives the long-distance flow of resources. The systemic distribution of photosynthate is known as assimilate partitioning and it is a major determinant of plant growth and productivity [7]. Our understanding of assimilate partitioning has advanced considerably over the last 30 years with the successful biochemical and molecular descriptions of several proteins that participate in this essential process (e.g. [8-11]). The current concept of phloem transport comprises three steps: (i) loading of photosynthates into the sieve element companion cell complex (se-cc complex) of minor veins in exporting leaves, (ii) translocation from source to sink, and (iii) unloading in growing or storing sinks [12].

Active transport by specific carriers across the apoplast, and symplastic transport via plasmodesmata, has been discussed as possible mechanisms for sucrose transport [13]. The transport is active and has been described as a sucrose-proton co-transport with a 1:1 stoichiometry [14].

Sink or source regulated modification of sucrose partitioning in plants is speculated to be a good strategy either for enhancing yield performance and improving plant-stress interactions, and for unravelling the biochemical, physiological and molecular mechanisms underlying sucrose partitioning in plants. To this end, modification could be achieved in two ways: (i) external treatments such as leaf girdling by hot wax collars to prevent export of assimilates from the leaves [15] or such as defoliating [16], or (ii), in vivo molecular manipulation [17,18]. In regard to the latter, one of the molecular

candidates for increasing or decreasing sink and source strength through intervention with assimilate loading or unloading is sucrose transporters. Another powerful tool for this is

the

15. Goldschmidt EE, Huber SC (1992) Regulation of photosynthesis by end-product accumulation in leaves of plants storing starch, sucrose, and hexose sugars. *Plant Physiol* 99: 1443-1448
16. Lee JM, Donaghy DJ, Roche JR (2008) Effect of defoliation severity on regrowth and nutritive value of perennial ryegrass dominant swards. *Agron J* 100:308-314
17. Lemoine R (2000) Sucrose transporters in plants: update on function and structure. *Biochim Biophys Acta* 1465: 246-262
18. Leggewie G, Kolbe A, Lemoine R, Roessner U, Lytovchenko A, et al. (2003) Overexpression of the sucrose transporter *SoSUT1* in potato results in alterations in leaf carbon partitioning and in tuber metabolism but has little impact on tuber morphology. *Planta* 217: 158-167.
19. Van Camp W (2005) Yield enhancement genes: seeds for growth. *Curr Opin Biotechnol* 16: 147-153
20. Sonnewald U1, Brauer M, von Schaewen A, Stitt M, Willmitzer L (1991) Transgenic tobacco plants expressing yeast-derived invertase in either the cytosol, vacuole or apoplast: a powerful tool for studying sucrose metabolism and sink/source interactions. *Plant J* 95-106
21. Koch KE, Wu Y, Xu J (1996) Sugar and metabolic regulation of genes for sucrose metabolism: potential influence of maize sucrose synthase and soluble invertase responses on carbon partitioning and sugar sensing. *J Exp Bot* 47:1179-1185