5BSHFUJOH 4NPLJOH $FTTBUJPO BOE 8FJHIU -PT "DDFQUBODF BOE $PNNJUNFOU 5IFSBQZ "$5 "Q

Jayson J Spas [1]*, J S Rossi [2], N D Collette [1]

[1]Rhode Island College, Department of Psychology, United States Providence, Rhode Island, United States
[2]University of Rhode Island (URI), Department of Psychology, United States Providence, Rhode Island, United States

Introduction

Smoking and obesity are the rst and second leading causes of preventable deaths in the United States (U.S.) [1]. With approximately 20% [2,3] of adults smoking and 36% and 34% of adults meeting criteria for overweight and obesity, smoking and excess weight (i.e., overweight and obesity) are among the most pressing health issues in the U.S. [4,5]. Along with premature death and disability, smoking and excess weight are associated with increased risk for acute medical events including myocardial infarction (MI) and cerebrovascular accident (CVA) as well as chronic disease including several cancers, cardiovascular disease (CVD), hypertension, hyperlipidemia and type 2 diabetes mellitus (T2DM). Beyond mortality and morbidity, the short- and long-term e ects of smoking and excess weight have staggering economic consequences. For example, the direct cost of smoking and obesity on the healthcare system is $190 and $193 billion per year, respectively, with obesity recently passing smoking as the greatest expense [6]. Furthermore, the combined cost of smoking and excess weight on the public and private sectors is remarkable, comprising more than 30% of total healthcare costs and hundreds of billions in lost productivity annually [7,8]. Overall, both smoking and excess weight are harmful patterns of consumptive behavior that persist despite serious physical, medical and economic consequences.

Standard intervention for smoking cessation and weight loss is either behavioral, pharmacological or a combination of both [9,10]. Interventions not uncommon for smoking cessation include quitting “cold turkey” and computer-tailored interventions (i.e., behavioral), the use of Varenicline or nicotine replacement therapy (NRT) such as the patch or lozenges (i.e., pharmacological) or some combination. Similarly, interventions not uncommon for weight loss include caloric restriction and increased physical activity (i.e., behavioral), the use of weight loss pills or other dietary supplements to suppress appetite and/or inhibit the absorption of fat during digestion (i.e., pharmacological)

diagnosis and treatment. Journal of Consulting and Clinical Psychology 64: 1152-1168.

27. Hayes SC, Lillis J (2011) Acceptance and Commitment Therapy; American Psychological Association.

28. Pennebaker JW (1997) Writing about emotional experiences as a therapeutic process. Psychological Science 8: 162-166.

29. Hayes SC (1987) A contextual approach to therapeutic change; in: Jacobson N, (ed): Psychotherapists in Clinical Practice. New York, Guilford Press 327-387.

30. Hayes SC, Wilson KG, Gifford EV, Bissett R, Batten S, et al. (2004)