
***Corresponding author:** Appakalai N Balamurugan, Department of Surgery, Clinical Islet Cell Laboratory, Centre for Cellular Transplantation, Cardiovascular Innovation Institute, University of Louisville, Louisville, KY, USA, Tel: 502-852-1131; Fax: 502-852-1391; E-mail: Bala.appakalai@louisville.edu

Received December 16, 2016; **Accepted** January 30, 2017; **Published** February 06, 2017

Citation: Dhanasekaran M, Loganathan G



to successfully adapt it for each individual organ. It is not as simple as it seems while coming across as a seemingly simple procedure; IAT requires highly specialized and delicate techniques in order to obtain an adequate number of healthy islets from a variety of donor types [5]. This report aims to describe the technical aspects of pancreas processing and islet isolation that are critical for achieving successful islet auto-transplantations.

Human islet isolation is a time sensitive and a skilled procedure performed by a well-trained team of members led by experienced personnel. Every team member's role should be carefully allocated to ensure an efficient and an effective way to isolating healthy islet for transplants. The islet laboratory should be adequately prepared prior to the isolation process. Using aseptic precautions, setting up of the biological safety cabinets (or laminar flow hood) with necessary materials for pancreatic trimming, cannulation and distension, digestion, recombination, purification and transplant bag preparation. On the other hand the remaining members should prepare the media and other in-use solutions that will be needed during the isolation. All necessary instruments such as centrifuges and thermal probes should be timely validated and turned on so that there would be no delay once after the pancreas has arrived.

For autologous isolations, the pancreas is dissected and immersed immediately in cold preservation solution. Generally the excess fat, connective and duodenal tissues are removed before packing on ice [6,7]. Cold storage preservation relies on hypothermia and carefully tailored solutions to slow metabolism, inhibit endogenous enzyme activity and support critical cellular processes despite the loss of an oxygenated blood supply. Organ packaging methods and solution ingredients have been designed to address several key problems associated with hypothermic ischemia followed by reperfusion including cellular swelling, ionic imbalance, acidosis, calcium accumulation and the production of reactive oxygen species. University of Wisconsin (UW) solution was developed in 1986, specifically for pancreas cold storage preservation [8]. UW contains phosphate, large molecules like saccharide ramoside, anionic lactobionate, allopurinol, glutathione, adenosine and a high, intracellular-mimicking K^+/Na^+ ratio [9]. While UW has become the standard organ transport solution, it is also costly with a short shelf-life and many of the ingredients, designed to inhibit tissue degradation, interfere with the catabolic activity of collagenase and neutral protease [10]. Other cold storage solutions have been proposed including Histidine-Tryptophan-Ketoglutarate (HTK), Celsior and the Kyoto solutions but UW remains the most common for pancreas hypothermic preservation. In trimming solution, a modified UW reverses the Na^+/K^+ ratio to mimic the natural extracellular environment and exchanges lactobionate for the less expensive but equally effective gluconate [11].



manual. The more effective the enzyme distention, the lesser the mass of undigested tissue and better is the islet yield. Historically, enzyme solution was loaded directly into the ductal cannula with a hand-held syringe, relying on retrograde perfusion to distend the pancreas [21]. In addition to improved distension and yield, automated pump perfusion provides precise control over injection pressure and enzyme solution temperature. The modern automated perfusion system is equipped with peristaltic pumps, two pressure sensors, a heater, a touch-screen, and data acquisition software (Bio-rep) that combines the convenience of hands-free automation with the flexibility to make manual adjustments to a variety of programmable parameters.

Distension pressure, pump speed, flow rate and temperature can all be monitored and controlled using a semi-automated perfusion system. Throughout the enzyme perfusion process, the temperature is kept between 6 and 16°C while the desired perfusion pressure is maintained between 60 and 80 mm Hg for the first 4 min, and gradually increased to 160-180 mm Hg until completion (approximately 10-12 min total distention time). However, perfusion pressure can vary significantly depending on the condition of the organ. Distention pressures could be low for a severely damaged, leaking, pancreas or high for an organ with abnormal ductal anatomy





additional 100 mL of media as a rinse solution. If the patient has no known allergy to ciprofloxacin, add 0.4 mL of Cipro[®] (1%=10000 µg/mL) to each volume of rinse media.

After attaching a 60 mL syringe to the transplant bag, place the syringe upright in a clamp stand and transfer the 100 mL tissue suspension into the bag through the syringe. Rinse the tissue conical twice with 50 mL volumes of rinse solution to transfer any residual islets. Aseptically recap and clamp the bag's inlet tubing to ensure a thorough seal for transport. The sealed bag should be gently rocked to evenly suspend the islets. Repeat these steps for additional transplant bags if needed. Once the transplant physician at the operation room has been alerted, the islet preparation can be readied for transport in a room temperature cooler equipped with temperature stabilizers [40].

Autologous islet isolation and transplantation has repeatedly demonstrated the ability to improve clinical outcomes by diminishing the impact of iatrogenic diabetes on patients undergoing pancreatectomy to alleviate CP or other disabling conditions. As practical experience has accumulated at an increasing number of qualified isolation centers, islet yield and viability, critical factors for achieving post-operative insulin independence, have progressively improved. It is imperative to understand and improve the technical aspects of the isolation procedure like the introduction of the simplified ATGS to improve purification yield [34] and the identification of post-isolation factors that detriment graft function [34]. Our research focus is on the mechanics of enzyme digestion, proposing a new enzyme mixture and variable dose classes that have increased the flexibility of the procedure to respond to different donor and tissue characteristics [17,20]. Despite these and other advances, the islet yield is significantly less than the available stores, indicating the need for more studies on efficient digestion. This demands a better understanding of the enzyme mechanics-Collagenase vs. neutral protease, their functional ingredients and interactions with different ECM components. This also necessitates the need for more specific techniques to overcome these inevitable obstacles. Furthermore, there is currently a heavy cost burden to establish facilities and perform these procedures, which severely limits their availability, especially in developing countries. All such technical and socio-economic parameters must be taken into consideration in order to successfully further develop and improve islet yield and transplantation outcomes for patients with CP.

Acknowledgment

The authors thank the Jewish Heritage Fund for Excellence for providing generous support to our program. The authors sincerely thank Kentucky Organ Donor

Kleinert and Brian Gettler for their assistance.

References

1. Majumder S, Chari ST (2016) Chronic pancreatitis. *Lancet* 387: 1957-1966.
- 2.



30. Balamurugan AN, Chang Y, Fung JJ, Trucco M, Bottino R (2003) Flexible management of enzymatic digestion improves human islet isolation outcome from sub-optimal donor pancreata. *Am J Transplant* 3:1135-1142.
31. Tsukada M, Saito T, Ise K, Kenjo A, Kimura T, et al. (2012) A Model to Evaluate Serine Protease Inhibitor in the Protection of Islets. *Cell Transplant* 21: 473-482.
32. Loganathan G, Graham ML, Radosevich DM, Soltani SM, Tiwari M, et al. (2013) Factors affecting transplant outcomes in diabetic nude mice receiving human, porcine, and nonhuman primate islets: analysis of 335 transplantations. *Transplant* 95: 1439-1447.
33. Lake SP, Bassett PD, Larkins A, Revell J, Walczak K, et al. (1989) Large-scale 2991 cell separator. *Diabetes* 38: 143-145.
34. Anazawa T, Matsumoto S, Yonekawa Y, Loganathan G, Wilhelm JJ, et al. (2011) Prediction of pancreatic tissue densities by an analytical test gradient system allotransplantation. *Transplant* 91: 508-514.
35. necessary? *Am J Surg* 166: 538-542.
36. cytokine chemokine production from human islet preparations, leading to prolonged beta-cell survival during pretransplantation culture. *Transplant Proc* 41: 314-315.
37. Chadwick DR, Robertson GS, Rose S, Contractor H, James RF, et al. (1993) UW solution and the roles of its individual components. *Transplant* 56: 288-293.
38. London NJ, Swift SM, Clayton HA (1998) Isolation, culture and functional evaluation of islets of Langerhans. *Diabetes Metab* 24: 200-207.
39. *Transplant* 5: 1-2.
40. Kaddis JS, Hanson MS, Cravens J, Qian D, Olack B, et al. (2013) Standardized transportation of human islets: an islet cell resource center study of more than 2,000 shipments. *Cell Transplant* 22: 1101-1111.

Citation: Dhanasekaran M, Loganathan G, Narayanan S, Tucker W, Jawahar AP, et al. Pancreatitis Pancreas (CPP) for Islet Auto-Transplantation (IAT). *J Clin Exp Transplant* 1: 112.

U U

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

U

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>