

The First Genetic Map and Positions of Major Fruit Trait Loci of Bitter Melon (*Momordica charantia*)

Chittaranjan Kole^{1*}, Bode A. Olukolu², Phullara Kole¹, Virendra K. Rao³, Anju Bajpai⁴, S. Backiyarani⁵, Jogendra Singh⁶, R. Elanchezhian⁷ and Albert G. Abbott¹

D. a. G. B. J. Ha. C U. C USA

Abstract

Cndgkv"gzvypukxg"ewnvkxcvkqp"qh"dkvgt"o gnqp"dqv j"cu"xg i gvcdbg"cpf" medicine in many countries of Asia, Africa, and South America, pq"ugtkqwu" g qtvu" jcxg" dggp" ocfg"hqt" igpgvke" cpf" dtggfkpi" uvwfkgu"qp"vjku" :qtrjcpø"etqr0"kp"eqpvtcu"vq" rqrwnct"ewewtdkvu." kv"ncemu" c" igpgvke"nkpmc i g" o cr"cu"tgswtgf"hqt" igpq o ke" fgrkevqp" and precise breeding. We report here on the construction of vjg"Łtuv" i gpgvke"nkpmc i g" o cr"qh"dkvgt" o gnqp"wukpi" c"ugv"qh"368" H4" rtqigpkgu" fgkxgf" htqo" cp" kpvgt/dqvcpkcn" xctkv{" etquu" dgvyggp" Vkycp" Y jkvg." Oqoqtfkcc" ejctcpvk" xct0" ejctcpvk." cpf" EDO34." O0"ejctcpvk" xct0" o wtkevc0" Vjku" o cr" eqpukuvu" qh' 32: "CHNR" o ctmgtu"cpf"Łxg"swcnkvcvkxg"vtckv"nqek"fkurgtugf"qxgt" 33"nkpmc i g" itqwrurcppkpi" c"vqvcn"fkuvcepg"qh"528209"eO0"Vjg"

xg"/

rgr

gr gf" v8

v8

Vjg"