

	Cycle	Temperature	Time
Ú{^Êâ^}æ~!æó[]		J1°Ô	I {i}
		J1°Ô	Fí •
V[~&Êâ[,) ÚÓÛ	Fí &^& ^•	Î€°Ô	Fí • Ç=ÊÊËÏ
		ÏG°Ô	H€ •
		J1°Ô	Fí •
U!âi}æi^ ÚÓÛ	Fí &^& ^•	IJEÍ°Ô	Fí •
		ÏG°Ô	H€ •
Òçc^}•i[]		ÏG°Ô	GE {i}
Ô[]•^!çæó[]		I°Ô	

ŠVHFÍ	Í	IÉĠİF	FÉİFİG	€İÍ	€İİİJ	€İG	€İİİG
ŠVHİF	Í	GÉFGİİ	€ÉJİH	€FÍ	€ÉHİİ	€İG€Í	€ÉH
ŠVIFÍ	H	GÉİİ	€ÉJİF	€I	€ÉFİİ	€HÍFÍ	€É€JF
ŠVİİÍ	H	FÉIHİJ	€ÉİÍ	€F	€ÉHÉJ	€İİİG	€ÉHÉIF
ŠVİİİ	I	H€Éİİ	FÉFJİG	€ÉHİİ	€ÉİHJ	F€€ÉİI	€ÉİIJ
ŠVJGF	I	GÉJGÍ	FÉFÍFÍ	€ÉİHG	€Éİİ	F€€FİIG	€ÉİİG
ŠVJİİ	I	HÉFGİF	FÉGEGH	€HÍ	€ÉİİJ	€ÉIJFJ	€ÉİEH
T^æ}	IÉHÉİİ	GÉJİFÍ	FÉFİG	€ÉHJİ	€ÉİİF	€Hİİ	€Éİİ

Table 4 Genetic diversity of 26 SSR loci within 5 provenances of *L. tulipifera* Note: An average number of alleles, Ne 9 Wj Y




Figure 2 Dendrogram of *L. tulipifera* in 5 provenances

Discussion

Genetic diversity of five *L. tulipifera* provenances