5IF 1IZTJDBM 5IFSBQZ "TTFTTNFOU BOE .BOBH $POHFOJUBM .VTDVMBS 5PSUJDPMMJT " 4VSW 1SPUPDPM GPS $.5

Anna M Öhman [1]*, Eva-Lott Mårdbrink [2], Christina Orefelt [3], Anthea Seager [4], Lisen Tell [5] and Eva Arrenhius Klackenberg [6]

[1,2]Department of Paediatrics, University of Gothenburg, Queen Silvia Children's Hospital, Gothenburg, Sweden
[3,5]Sollevi Physiotherapy, Stockholm, Sweden
[4]Senior Physiotherapist,Temple Street Children's University Hospital, Dublin 1, Ireland
[6]Department of physiotherapy, The Astrid Lindgren Children´s Hospital, Stockholm, Sweden

Abstract

Background: Infants with congenital muscular torticollis (CMT) are treated with physical therapy, however the knowledge about type of assessment and treatment in current clinical practice is not clear.

Aim: This study aimed to investigate the management of infants with CMT within a network of physical therapists.

Settings/population: Physical therapists from a network for CMT participated in the study. Method: With permission a questionnaire from New Zealand was used. Subsequently an expert group of physical therapists developed a draft assessment protocol.

Result: In general there was a high degree of consensus between the respondents in the current survey. For the assessment visual estimates were most commonly used and an evaluation of cervical muscle strength of the neck was always/often conducted. The most effective form of intervention as perceived by the respondents in the management of CMT was passive stretching, handling advice, facilitation with strengthening exercises of the neck muscle and facilitation of active cervical range of motion. There were big discrepancies in assessment of hand- and hip asymmetry.

Conclusion: There seems to be a need to develop international guidelines for CMT to improve the evidence practice of assessment and treatment.

Keywords: Congenital Muscular Torticollis, Physical erapy, Assessment, Treatment

Introduction

Congenital muscular torticollis (CMT) is the third most common musculoskeletal abnormality in infants next to hip dysplasia and clubfoot. e reported incidence is 0.4-2.0% [1,2] however a recent study indicates that it might be higher [3]. CMT is a result of shortening or excessive contraction of the sternocleidomastoid (SCM) muscle with limited range of motion (ROM) in both rotation and lateral exion of the neck and an imbalance of muscle function around the neck [4-6]. Due to the positional preference there is a high risk that infants with CMT develop deformational plagiocephaly [7-9]. e birth history demonstrates an unusually high incidence of di culties during labour e.g. breech presentation is commonly found [10-14]. ere is also a coexistence with hip dysplasia [10,11,14]. Infants with CMT are found to be at risk of delay in achieving motor milestones [15,16].

Evidence-based Physical therapy is important for patients because it implies that they will be o ered the safest and most e ective interventions within the limitations of current knowledge. e expectation is that this will produce the best possible clinical outcome [17]. Physical therapists in general have positive attitudes and beliefs regarding evidence-based practice (EBP) [18,19]. However, the majority of physical therapists indicated that they need to increase the use of evidence in their daily practice [18]. e process of implementing any research outcome begins with awareness [20]. In order to understand to what extent there is a need to improve and implement "new" methods and strategies for assessment and treatment of CMT among physical therapists we need to know more about the current practice. Luxford et al. investigated current physical therapy management and issues in management of infants with CMT in New Zealand (NZ) [21]. ey constructed a questionnaire for this purpose. In their survey they found that most physical therapists in NZ used a visual estimate [21], this may re ect a lack of knowledge about measurement tools or a possible reluctance to change. Luxford et al. [21] identi ed key points such as consideration of the use of international protocols, routine assessments of infants hips and the use of information handouts to parents. e aims of physical therapy treatment for infants with CMT are to prevent/reduce plagiocephaly and to achieve symmetrical head position, ROM and muscle function of the neck. It is also important to be observant on the infant's motor development. Assessments and treatments are described in several studies [1,5,22,23].

e aim of the current study is to obtain knowledge about the existing practice in the physical therapy management of infants with CMT among members of a network for torticollis and also to develop a dra assessment protocol that could be used when examining an infant with CMT. e rst author is responsible for the survey.

Method

Eighty-nine members of the network for physical therapists working with torticollis were invited to participate in this survey by

Received May 09, 2013; Accepted June 28, 2013; Published June 30, 2013

Citation: Öhman AM, Mårdbrink EL, Orefelt C, Seager A, Tell L, et al. (2013) The Physical Therapy Assessment and Management of Infants with Congenital Muscular Torticollis. A survey and a Suggested Assessment Protocol for CMT. J Nov Physiother 3: 165. doi:10.4172/2165-7025.1000165

Copyright: © 2013 Öhman AM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted

e-mail contact. Most of the members are from Sweden but a few are from Denmark. Forty physical therapists (45%) chose to participate. With permission from Bernadette Luxford, the NZ questionnaire was used. It was translated to Swedish, however the participants had access to both the English and the Swedish version if they wanted.

e questionnaire is anonymous, 10-pages long and comprises 28 questions divided into ve sections collecting general and demographic information, information speci c to subjective assessment, objective assessment techniques, di erential diagnosis and treatment techniques used in the management of CMT [21]. e only di erence in the Swedish version is that one assessment tool is added in one of the questions. In addition to the arthrodial protracto another large protractor is some-times used in Sweden [22,24].

At the beginning of the questionnaire there is a screening question to ensure that the information is sought from physical therapists that had assessed and/or treated an infant (under two years of age) with CMT within the last year. Descriptive statistic is used for the result.

A er the survey, a panel of experts developed a dra assessment protocol. e panel consisted of six pediatric physical therapists experienced in the assessment of infants with CMT. e panel members had professional experience of assessment of infants with CMT ranging from 10 to 40 years, with a mean of 18 years.

Result

Of the 40 physical therapists who replied three had not treated any infant during the last year, therefore the analysis is done on the data from the remaining 37 respondents. irty-four physical therapists from twelve districts in Sweden and three physical therapists in Denmark were included; one participant did not indicate which region she was from. All participants were female, and 21 were aged between 30-49 years old. e 37 respondents had worked as physical therapists for median of 20 years (range 5-40 years), in pediatric settings for median of 14 years (range 1-40 years). ey assessed/treated in median 13 infants with CMT during a year, range 1-300 infants.

Nine sources of referral to physical therapy were identi ed; the three most common were “child health care centers” (81%), commonly referred to as “well baby clinics”, pediatricians (73%), and “special units for children” (46%). Eighty-four percent of the respondents had an information handout for parents. Sixty-two percent had best practice guidelines for CMT at their place of employment and 16% reported that a guideline was in the process of being developed.

When considering the subjective assessment there is a high level of agreement about the information sought (Table 1). With regard to the objective assessment techniques for infants with CMT there is also high agreement about most components (Table 2). For hand

convenient to use, there may be some reluctance to change. ROM in infants is not very easy to assess, as the infants dislike to be held during measurements.

Treatment

e most common treatments were passive stretch, handling advice, facilitation of neck muscle strength and facilitation of active

head