**Keywords:** 1- k., -., :C., :H., ..., :E.; H., ..., :E.; E.;

#### Introduction

 وها أن أن المرسم و عدد الله يعون مع مراح مع مواهم و عدد الله مع والم عدد و المراح و الله الله و المراح و الله ا الأمام عدد منا و عمل مع مواهم و المعلق على مهامة المراح و المراح و الله المراح و الله المراح و الله المراح و ا مع المراح و الله الله الله المراح و الله المراح و المراح و الله المراح و المراح و الله المراح و المراح و الله المراح و المراح و الله المراح و المر

## **Conclusion**

. C<sub>سه</sub> ، سود والهر لهرو لهره بردو وريس و و .

### Acknowledgment



# Con ict of Interest



#### References

- Palmer BW, Heaton SC, Jeste DV (1999) Older patients with schizophrenia: challenges in the coming decades. Psychiatric Services 50: 1178-1183.
- Patorno E, Bohn R, Wahl P, Avorn J, Patrick AR, et al. (2010) Anticonvulsant medications and the risk of suicide, attempted suicide, or violent death.JAMA 303: 1401-1409.
- Olesen JB, Hansen PR, Erdal J, Abildstrøm SZ, Weeke P, et al. (2010) Antiepileptic drugs and risk of suicide: a nationwide study. Pharmacoepidem Dr S 19: 518-524.
- Leipzig R, Cumming R, Tinetti M (1999) Drugs and falls in older people: a systematic review and meta-analysis: I. Psychotropic drugs. J Am Geriatr Soc 47: 30-39.
- Gill S, Bronskill S, Normand S, Anderson GM, Sykora K, et al. (2007) Antipsychotic drug use and mortality in older adults with dementia. Ann Intern Med 146: 775-786.
- Casey D, Haupt D, Newcomer J, Henderson DC, Sernyak MJ, et al. (2004) Antipsychotic-induced weight gain and metabolic abnormalities: implications for increased mortality in patients with schizophrenia. J Clin Psychiatry 65(Suppl 7): 4–18.
- Schneider LS, Dagerman KS, Insel P (2005) Risk of Death with Atypical Antipsychotic Drug Treatment for Dementia. JAMA 294: 1934-1943.
- Meijer WEE, Heerdink ER, Nolen WA, Herings RMC, Leufkens HGM, et al. (2004) Association of Risk of Abnormal Bleeding With Degree of Serotonin Reuptake Inhibition by Antidepressants. Arch Intern Med 164: 2367-2370.