The silent epidemic: Understanding the health consequences of chronic stress

Nida Ali*

Department for Clinical and Health Psychology, University of Vienna, Austria

ABSTRACT:

KEYWORDS: Stress Hormones, Sleep Disturbances, Behavioral Changes.

INTRODUCTION

Chronic stress triggers a cascade of physiological responses in the body, primarily through the activation of the Hypothalamic-Pituitary-Adrenal (HPA) axis. This leads to the release of stress hormones, such as cortisol and adrenaline. While these hormones are essential for short-term responses to stress, their rgtukuvgpv"gngxcvkqp"ecp" j cxg" f gvtk o gpvcn"g gevu"*Dctvqnq o week" A, 1998). Prolonged exposure to stress hormones can lead to increased heart rate and blood pressure, heightening the risk of cardiovascular diseases, including hypertension, heart attack, and stroke. Chronic stress can suppress the immune system, making individuals more susceptible to infections and chronic diseases. Elevated cortisol levels can inhibit the production of cytokines, essential for immune response, thus k o rektkpi "vjg"dqf {øu"cdknkv { "vq"Łijv"q "knnpguugu"*Dqqpuvtc"T." 1945). Stress often leads to unhealthy coping mechanisms, such as overeating or poor dietary choices. This can result in weight gain and associated conditions like obesity, diabetes, and metabolic syndrome (Chandola T, 2006). Cortisol also promotes the accumulation of fat, particularly in the abdominal area. Chronic stress can exacerbate gastrointestinal disorders, including Irritable Bowel Syndrome (IBS), Gastroesophageal Tg ł wz "Fkugcug" * I GTF+. "cpf" wnegtu0" Uvtguu "c gevu" i wv "o qvknkv { " cpf"ecp"ngcf"vq"kpłcoocvkqp. "fkutwrvkpi"vjg"fgnkecvg"dcncpeg"qh" gut microbiota (Checkley S, 1996).

PSYCHOLOGICAL EFFECTS OF CHRONIC STRESS: Vjg"ru{ejqnqikecn"tcokŁecvkqpu"qh"ejtqpke"uvtguu" can be equally severe, impacting mental health and overall

Received: 02-Sep-2024, Manuscript No: ijemhhr-24-151444;

Editor assigned: 03-Sep-2024, Pre QC No. ijemhhr-24-151444 (PQ);

Reviewed: 17-Sep-2024, QC No. ijemhhr-24-151444;

Revised: 23-Sep-2024, Manuscript No. ijemhhr-24-151444(R); **Published:** 30-Sep-2024, DOI: 10.4172/1522-4821.1000656

*Correspondence regarding this article should be directed to: nidali@univie.ac.at

swcnkv {"qh"nkhg0" Ej tqpke"uvtguu"ku"c"uk i pkŁecpv"tkum"hcevqt"hqt" developing anxiety disorders and depression. Persistent stress can alter brain chemistry and function, leading to feelings of hopelessness and emotional instability (Conrad CD, 2010). Prolonged stress can impair cognitive functions, including memory, attention, and decision-making. The hippocampus, a region of the brain involved in learning and memory, can shrink under chronic stress, leading to fk ewnvkgu" kp" rtqeguukpi" kphqtocvkqp" cpf" hqtokpi" pgy" memories (Davidson LM,1986). Individuals experiencing chronic stress often face sleep problems, such as insomnia or disrupted sleep patterns. Poor sleep can further exacerbate stress and lead to a vicious cycle that deteriorates both physical and mental health (Marin MF, 2011).

Addressing chronic stress requires a multi-faceted approach that includes lifestyle changes, psychological interventions, and, when necessary, medical treatment. Practices such as mindfulness meditation, yoga, and deep-breathing gzgtekugu" ecp" jgnr" kpfkxkfwcnu" ocpcig" uvtguu" g gevkxgn{0" These techniques promote relaxation and can reduce the physiological impacts of stress. Regular exercise is a powerful stress reliever (McGonagle KA,1990). Physical activity increases the production of endorphins, which can elevate mood and improve overall well-being. A balanced diet rich in fruits, vegetables, whole grains, and lean proteins ecp" jgnr" o kvk i cvg"v jg"g gevu"qh"uvtguu0"Q o gic/5"hcvv{"cekfu." $hqwpf"kp" \verb"Luj" cpf" pwvu." \verb"jcxg" dggp" u \verb"jqyp" vq" uwrrqtv" dtckp"$ health and reduce symptoms of depression. Building a strong support network is vital for coping with chronic stress. Engaging with friends, family, or support groups can provide emotional support and practical assistance. For individuals struggling to manage chronic stress, seeking vjg"jgnr"qh"c"ogpvcn"jgcnvj"rtqhguukqpcn"ecp"dg"dgpgŁekcn0" Therapy, such as Cognitive-Behavioral Therapy (CBT), can equip individuals with coping strategies and tools to address uvtguuqtu"g gevkxgn{"*Uejokfv"OX.422:+0

Chronic stress cap "ngcf" vq" ukipkLecpv" dgjcxkqtcn" ejcpigu" that further impact mental health. Individuals may resort to unhealthy coping mechanisms such as substance abuse, overeating, or social withdrawal as a way to escape their stressors. These behaviors can create a feedback loop, exacerbating feelings of anxiety and depression while also isolating individuals from supportive social networks. Chronic stress can also manifest in irritability and mood uykpiu." ocmkpi" kv" fk ewnv" hqt" kpfkxkfwcnu" vq" ockpvckp" relationships and engage in daily activities. Over time, vjgug" ru{ejqnqikecn" g gevu" ecp" ukipkŁecpvn{" korckt" overall functioning, leading to diminished quality of life cpf" kpetgcugf" fk ewnv{" kp" o cpcikpi" dqvj" rgtuqpcn" cpf" professional responsibilities. Recognizing and addressing these psychological consequences is crucial for breaking the cycle of chronic stress and fostering healthier coping strategies (Tafet GE,2003).

CONCLUSION

Chronic stress is a sigpkLecpv"rwdnke"jgcnvj"eqpegtp"ykvj"hct/ tgcejkpi"g gevu"qp"rj{ukecn"cpf"ogpvcn"jgcnvj0"Tgeqipk|kpi" the signs and understanding the impacts of chronic stress can empower individuals to take proactive steps toward o cpcikpi" vjgkt" uvtguu" ngxgnu0" D{" korngogpvkpi" g gevkxg" coping strategies and seeking support when needed, kpfkxkfwcnu"ecp" o kvki cvg"v jg"cfxgtug"g gevu"qh"e jronic stress and enhance their overall health and well-being.

References

Bartolomucci, A., Palanza, P., Sacerdote, P., Panerai, AE., Sgoifo, A (1998) Social factors and individual vulnerability to chronic stress exposure. Neurosci Biobehav Rev.1;29(1):67-81.

Boonstra, R (1945). Reality as the leading cause of stress: rethinking the impact of chronic stress in nature. Funct Ecol. :27(1):11-23.

Chandola, T., Brunner, E., Marmot, M (2006). Chronic stress at work and the metabolic syndrome: prospective study. 332(7540):521-5.

Checkley, S (1996). The neuroendocrinology of depression and chronic stress. Br Med Bull. 1;52(3):597-617.

Conrad, CD (2010). C"etkvkecn"tgxkgy"qh"ejtqpke"uvtguu"g gevu" on spatial learning and memory. Neuro Psycho Bio Psych. 30;34(5):742-55.

Davidson, LM., Baum, A (1986). Chronic stress and posttraumatic stress disorders. J Consult Clin Psychol. 54(3):303.

Marin, MF., Lord, C., Andrews, J., Juster, RP., Sindi, S (2011). Chronic stress, cognitive functioning and mental health. Neurobiol Learn. Mem. 1:96(4):583-95.

McGonagle, KA., Kessler, RC (1990). Chronic stress, acute stress, and depressive symptoms. Amer J Comm Psych. 18(5):681-706.

Schmidt, MV., Sterlemann, V., Müller, MB (2008). Chronic stress and individual vulnerability. Ann Acad Sci. 1148(1):174-83.

Tafet, GE., Bernardini, R (2003). Psychoneuroendocrinological links between chronic stress and depression. Prog. Neuro Psycho Bio Psych. 1;27(6):893-903.