

# 5 I F S N P H S B Q I J D 1 S P G J M F P G 7 P M M F Z C B M M 1 M 1 S F W F O U J P O

Berzosa C<sup>1,2\*</sup>, Mejías-Martínez M<sup>1,2</sup>, Valero-Campo C<sup>1,2</sup> and Bataller-Cervero AV<sup>1,2</sup>

<sup>1</sup>San Jorge University, Villanueva de Gállego Zaragoza, Spain

<sup>2</sup>ValorA Research Group, Faculty of Health Sciences, San Jorge University, Spain

Keywords: ~~Sample~~ ~~Evidence~~ ~~Individualized~~ ~~SWMHD~~ ~~GRL~~ ~~EDT~~ ~~Domestic~~ ~~Product~~ ~~Waste~~ ~~Hazardous~~ ~~Toxic~~ ~~Value~~ ~~Control~~ ~~Waste~~ ~~Management~~ ~~Health~~ ~~Risk~~ ~~Assessment~~ ~~Policy~~ ~~Regulation~~ ~~Standardization~~ ~~Guidelines~~ ~~Classification~~ ~~Labeling~~ ~~Packaging~~ ~~Waste~~ ~~Management~~ ~~Health~~ ~~Risk~~ ~~Assessment~~ ~~Policy~~ ~~Regulation~~ ~~Standardization~~ ~~Guidelines~~ ~~Classification~~ ~~Labeling~~ ~~Packaging~~

Anterior zones (pectoral zones) and posterior zones (right and left scapular waist and lumbar zone) were delimited and mean temperature of each zone were calculated.