Case Report Open Access

Total Colectomy with an Ileorectal Anastomosis: Perforation of Distal Portion of the Terminal Ileum

Rubén Gonzalo González[1*], Sara Regano Diez[1], Maria Antonia de Andrés Fuertes[1], José Luis Conty S[1], Adela Delgado Tapia[2] and Francisco Javier Gil Piedra[1]

[1]*Department of General Surgery, Hospital Laredo, Spain*

[2]*Department of Radiology, Hospital Laredo, Spain*

[*]**Corresponding author:** Rubén Gonzalo González, Department of General Surgery, Hospital Laredo, Av Human Rights, s/n, 39770 Laredo, Cantabria, Spain, Tel: 620 108 834; E-mail: rubsoygg@hotmail.com

Rec date: July 16, 2015, **Acc date:** August 13, 2015, **Pub date:** August 20, 2015

Copyright: © 2015 González RG, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Total colectomy followed by ileorectal anastomosis is an established operation that may be employed as a surgical solution for a variety of colonic diseases. Postoperative morbidity and mortality rates are generally regarded as low, and functional outcome is rated as good to excellent by most patients. The ileorectal anastomosis is usually latero-terminal and is constructed with staples. If the segment of terminal ileum extending past the site of the anastomosis is too long, it remains as an appendage, and this portion of bowel may fold or twist upon itself, thus creating an intermittent closed-loop obstruction of its tip, leading to localized perforation, sepsis, and eventual fistula formation. The case of one patient who developed significant complications arising from this portion of the bowel is reported.

Keywords: Total colectomy; Ileorectal anastomosis; Intestinal perforation; Mechanical suture

Total colectomy (TC) with ileorectal anastomosis (IRA) is a relatively frequent procedure today. Indications are well defined, functional results are compatible with an almost normal quality of life, and postoperative morbidity and mortality rates are low [1-5]. TC can be performed through laparotomy or laparoscopy; IRA can be manual or mechanic, end-to-end or side-to-end. When performing side-to-end mechanical IRA, the circular stapler head is first introduced through the end of the ileum, leaving 5 cm of terminal ileum distal to the anastomosis, and the circular stapler is introduced through the anus. The stapler is then assembled and the anastomosis is completed. This end of distal ileum may act as an appendix or blind loop if it is too long. We present the case of a patient with a serious complication stemming from this portion of distal ileum.

This 64-year-old male patient had no previous significant medical history. Previous surgeries included laparoscopic TC with mechanical side-to-end IRA 10 years earlier to treat a synchronous colonic adenocarcinoma located at the sigmoid and the transversal colon (T3N0M0). Currently, he continued periodic follow-up visits and annual rectal endoscopy carried out by the Digestive Department with no significant findings.

He presented to the emergency department with abdominal pain at the left hemiabdomen. He had been suffering from this pain for the last 15 days, and in the last 12 hours, it increased sharply. He presented with nausea without emesis. His last bowel movement 24 hours earlier was reported to be normal. He had a temperature (38ºC) and was conscious and oriented, and was well hydrated, with normal color. Cardiopulmonary auscultation was normal. The abdomen was distended and tympanic. Palpation at the left hemiabdomen was painfulbut there was no guarding. No masses or hernias are palpated. The hemogram revealed leukocytosis (14,000 leukocytes with 75% neutrophils) and an increased C-reactive protein (10.70 mg/dL). Renal function, pancreatic enzymes, and hepatic profile were normal. Simple thoracic X-ray demonstrated a significant pneumoperitoneum (Figure 1). An urgent abdominal CT scan was performed. It showed the presence of a remarkable pneumoperitoneum at the intra-abdominal level, and a small bowel loop on the left side with a thickened/ inflammatory appearance with significant fraying of peripheral fat and a minimal amount of adjacent liquid (corresponding to the terminal ileum segment distal to the IRA). The possibility of this process being the origin of the current pneumoperitoneum could not be excluded.



Figure 1: Thoracic X-ray: Arrows point at the pneumoperitoneum.

At the pelvic level, the IRA was seen without complications (Figures 2-4). The decision was made to perform a median laparotomy which revealed purulent peritonitis secondary to a perforation on the border of the terminal ileum distal to the IRA. This approximately 15-cm-long small bowel segment (as an appendix or blind loop) was

hypertrophic, thickened, and dilated. No additional pathological findings were seen at the IRA or in the rest of the abdominal cavity.



Figure 2: The arrow points at the IRA. No complications seen.