

Treatment Reception in a Cross-sectional Study of US Adults with Communication Disorders

Nazanin Jagadapillai*

Department of Pediatrics, Pediatric Research Institute, University of Louisville, USA

Abstract

Communication disorders significantly impact the lives of adults in the United States, yet the receipt of treatment among a fected individuals remains inadequately understood. This study examines treatment reception among a nationally representative sample of US adults with communication disorders, utilizing data from a cross-sectional survey. Participants were assessed for the presence of communication disorders and queried on their utilization of treatment modalities such as speech therapy, medical interventions, assistive devices, and alternative therapies. Findings indicate a notable proportion of adults with communication disorders do not receive treatment, with disparities observed across demographic factors including age, gender, race/ethnicity, and socioeconomic status. Common barriers to treatment access include fnancial constraints, lack of healthcare coverage, and geographical limitations. Addressing these disparitic

ReviewedPresquited.

No- jspt-24-142709; Revised: 25-July-2024, Manuscript No: jspt-24-142709 (R);

Published: 31-July-2024, DOI: 10.4172/2472-5005.1000252

Keywords: Citation: Nazarin J (2024) Treatment Reception in a Cross-sectional Study of US Adults with Communication Disorders. J Speech Pathol Ther 9: 252. Healthcare access; Disparities; Speech therapy; Cross-sectional study

Copyright: © 2024 Nazarin J. This is an open-access article distributed under the

Copyright: © 2024 Nazanin J. This is an open-access article distributed under the Introduction in the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and Communication disputes are created.

minority groups were less likely to receive treatment compared to younger adults and those from non-minority groups.

Barriers to care: Common barriers to accessing treatment included nancial constraints, lack of healthcare coverage, limited availability of specialized services, and geographical barriers (especially in rural areas). ese barriers disproportionately a ected underserved populations, contributing to disparities in treatment reception.

Impact of untreated communication disorders: Untreated communication disorders can signi cantly impact individuals' quality of life, a ecting social interactions, employment opportunities, academic achievement, and overall well-being. Addressing these disorders through timely and appropriate treatment is essential for improving outcomes and reducing long-term healthcare costs.

Implications for healthcare policy and practice

e ndings underscore the importance of developing targeted interventions to improve access to care for adults with communication disorders.

Enhancing healthcare access: Expanding insurance coverage for speech therapy and related services, particularly for vulnerable populations.

Workforce development: Increasing the number of trained healthcare professionals specializing in communication disorders, especially in underserved areas.

Telehealth: Promoting telehealth services to overcome geographical barriers and improve access to specialists.

Conclusion

e study highlights signi cant disparities in treatment reception among US adults with communication disorders and identies barriers that hinder access to care. Addressing these disparities requires a multifaceted approach involving healthcare policy reforms, workforce development, and innovative healthcare delivery models. By improving access to timely and appropriate treatment, healthcare providers and policymakers can enhance the quality of life for individuals with communication disorders and promote more inclusive and equitable healthcare practices. e ndings from this cross-sectional study provide valuable insights into the treatment reception among US adults with communication disorders, shedding light on both successes

and challenges within the current healthcare landscape. Our analysis revealed signi cant disparities in treatment access across demographic factors, including age, gender, race/ethnicity, and socioeconomic status. ese disparities underscore the need for targeted interventions to improve equity in healthcare delivery for individuals with