Journal of
Bioremediation & Biodegradation

Liu et al., J Bioremediat Biodegrad 2016, 7:4
DOI:10.4172/2155-6199.1000355

Research Article Open Acces

Abstract

Polycyclic Aromatic Hydrocarbons (PAHs) are widely distributed in the environment originating from both natural and anthropogenic sources. Consequently, they are found ubiquitously in the lithosphere, hydrosphere, atmosphere and biosphere. They are of major concern due to their recalcitrant nature, ability to bio-accumulate and their toxic, carcinogenic and mutagenic effects on biological systems. One major anthropogenic source of PAHs is creosote which is used as a wood preservative and contains 85% PAHs (w/w). Industrial facilities carrying out creosote treatment activities can result in major contamination of soil and groundwater at these sites. The current study investigates the potential of Ecopiling as a bioremediation strategy for creosote contaminated soil/sediment at a former wood preservation facility. The study involved greenhouse experiments to assess the phytoremediation potential of perennial ryegrass (Lolium perenne L.) and onsite construction of two small-scale, trial Ecopiles. The greenhouse study showed that the creosote contaminated sediment had an inhibitory effect on plant growth and UHGXFHG WKH HIIHFWLYHQHVV RI UKL]RUHPHGLDWLRQ +RZHYHU WKH XVH RI D F in the phyto-toxicity experienced by the ryegrass plants and subsequently resulted in greater PAH degradation in these pots. The use of Ecopiling was found to reduce the sum of EPA 16 PAH concentrations in the sediment by 94-96% after 730 days. The incorporation of spent-mushroom compost into the Ecopile, did initially increase the rate RI 3$+ GHJUDGDWLRQ KRZHYHU DIWHU GD\V WKHUH ZDV QR VLJQL¿FDQW GLI remaining in the two Ecopiles. Although Ecopiling is a slow process compared to thermal and chemical treatments it has proven to be an effective and low cost process for the remediation of PAHs from industrial sediments.

Keywords: Ecopiling; Bioremediation; Phytoremediation; PAHs; Field trials

Introduction

Polycyclic Aromatic Hydrocarbons (PAHs) are a group of compounds composed of fused aromatic rings. ere are 1896 possible PAH structures containing two to eight fused rings. Many PAHs occur in the environment as a result of natural combustion processes such as forest res and volcanic activity [1]. Anthropogenic inputs of PAHs into the environment stem from burning of fossils fuels, waste incineration, smelting operations and the use of oil and coal products such as asphalt and creosote. With the exception of leachate and spills, the vast majority of anthropogenic PAHs enter the environment in the form of air pollutants and over time are transported into water, soils, sediments and biota [2]. Low molecular weight PAHs (2-3 ring compounds) tend to be more water soluble than the higher molecular weight PAHs (5-8 rings) and these tend to partition between the vapour and solid phases. Heavier PAHs tend to be exclusively found in the solid phase and so are generally adsorbed onto dissolved organic matter (DOM) or soil particulates [3]. e tendency of PAHs to accumulate in soils, sediments and biota increases with increasing molecular weight of the PAH and the corresponding increase in the soil/sediment partition coe cient (K_{oc}). PAHs themselves are generally unreactive and have low acute toxicity. However, their metabolites can be potent mutagens and carcinogens [4].

Creosote is a mixture containing 85% PAHs (w/w), 10% phenolics and 5% heterocyclic compounds and is commonly used as a wood preservative to treat wooden structures such as rail-road sleepers, electricity line poles and wooden marine structures. It is a dense, non-aqueous phase liquid (DNAPL) that contaminates soil and water at many wood-preservation sites [5,6]. In the past, ine ciencies in the wood impregnation process led to losses of up to 2% of the creosote charge in the form of fugitive air emissions and drip loss to the underlying ground (with typical charges being 120 L of creosote per m^3 of wood). Remediation of these sites is di cult as aged PAHs tend to be tightly bound to soil particles reducing their bioavailability for microbial degradation. Soils and sediments with high clay content are particularly problematic due to the strong absorption and adsorption of PAHs to clay particles [7]. Currently, there are no European wide soil quality standards for PAHs in soil. However, the Dutch target and invention levels are widely used as benchmarks in remediation e orts.

Copyright: © 2016 Liu X, et al. This is an open-a ccess article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

over the 730 day period. e other remaining PAHs were degraded by 80-99%. e PAH analysis of the Ecopile soils suggested that there was considerably more degradation of phenanthrene, uoranthene, benzo[a]anthracene, benzo[a]pyrene and indeno[1,2,3,cd]pyrene in Ecopile 1 than in Ecopile 2. Many of the unidenti ed compounds present in the initial contaminated sediment (unidenti ed peaks) were not observed in the GC-FID pro les a er 161 days, suggesting that these compounds were also biodegraded (Figure 5). A control pile containing just untreated sediment showed a 27.5% reduction in EPA 16 PAH concentrations over the 730 day period.

Discussion

is study examined the feasibility of using Ecopiling as a possible remediation system for the removal of PAHs from creosote contaminated sediment. Ecopiling is a modi cation of traditional passive composting (biopiling) by incorporating phytoremediation into the remediation process. e Ecopile process also involves bio-stimulation of indigenous hydrocarbon degraders, bio-augmentation through inoculation with known hydrocarbon degrading consortia and phytoremediation, through the e ect of root growth and penetration

kg^{-1}). However, they reported up to a 50% decrease in root and shoot yield in ryegrass. e reduction in growth was thought to be due to toxicity of low molecular weight constituents. Yu et al. [20] showed that both phenanthrene and pyrene signi cantly decreased the biomass of ryegrass plants. While in an earlier report, Yu et al. [21] found no e ects on plant growth or any sign of stress in white clover and perennial ryegrass grown in soil spiked with phenanthrene and pyrene at rates up to 375 mg kg^{-1}. Saison et al. [22] found that the growth of Italian rye-grass (*Lolium multiflorum var. Fastuliform*) and red clover (*Trifolium pratense*) on coking plant soils with 2077 mg PAH kg^{-1} soil was similar to plant growth in an uncontaminated control. In the current study we found that sowing the plant seeds in a thin layer of clean soil led to signi cant improvements in plant biomass. Seed germination is a very sensitive phase of plant growth, and many classical toxicity assays are based upon germination or seedling emergence in the presence of contaminants. Henner et al. [23] found that seed germination was strongly inhibited by low molecular weight, water-soluble compounds, for example toluene, styrene and naphthalene. Kuiper et al. [24] also found that naphthalene reduced seedling emergence. Seedlings grown in naphthalene contaminated (200 mg kg^{-1}) soil produced no above ground growth. Liu et al. [25] found the high levels of phenanthrene (200 mg kg^{-1}) exhibited toxic e ects on ryegrass but this toxicity was only observed a er 75 days. e analysis of the soils in the current study showed that there were low molecular weights PAHs present which may have been responsible for the reduced biomass observed.

In the greenhouse trials the growth of ryegrass in the creosote contaminated soil did result in greater PAH degradation than in the unplanted controls. Glass beakers with greater ryegrass biomass had greater levels of PAH biodegradation. Yu et al. [21] showed that ryegrass enhanced the dissipation of phenanthrene and pyrene through the signi cant (p<0.05) increase in peroxidase activities in soil. Olson et al. [26] found that planting soil with perennial ryegrass led to 85% dissipation of PAHs in contaminated soil. ere are also many other reports of (s)5 -7 (59 (n45 (s)sw726 Tm [(<0.0)4 (e2 (r)-9.9 (ts oe 0 0 9 42.519n (e)13 (ep)-9 vps en)-5 (h)3 (a)9 (n)4 (ce)-5 (d t (

was characterised by having a high diversity and the persistence of an introduced bacterial consortium during the entire biotreatment process. Llado et al. [36] used a dynamic biopile as a remediation

25. /LX 6/ &DR =+ /LX +((IIHFW RI U\HJUDVV /ROLXP PXOWLÀRUXP / JURZWK on degradation of phenanthrene and enzyme activity in soil. Plant Soil Environ 59: 247-253.

26. Olson PE, Castro A, Joern M, Duteau NM, Pilon-Smits E, et al. (2008) Effects of agronomic practices on phytoremediation of an aged PAH-contaminated soil. - (QYLURQ 4XDO

27. Günther T, Dornberger U, Fritsche W (1996) Effects of ryegrass on biodegradation of hydrocarbons in soil. Chemosphere 33: 203-215.

28. Binet P, Portal J, Layval C (2000) Fate of polycyclic aromatic hydrocarbons in the rhizosphere and mycorrhizosphere of Ryegrass. Plant and Soil 227: 207-213.

29. Afzal M, Yousaf S, Reichenauer TG, Sessitsch A (2012) The inoculation method affects colonization and performance of bacterial inoculant strains in the phytoremediation of soil contaminated with diesel oil. Int J Phytoremediation 14: 35-47.

30. 7HQJ < 6KHQ < /XR < 6XQ ; 6XQ 0 HW DO ,QÀXHQFH RI 5KL]RELXP meliloti on phytoremediation of polycyclic aromatic hydrocarbons by alfalfa in an aged contaminated soil. J Hazard Mater 186: 1271-1276.

31. Pritchard PH, Jones-Meehan J, Straube W, Nestler C, Jones W, et al. (2002) Polycyclic aromatic hydrocarbons (PAHs): Improved land treatment with bioaugmentation. p. B1-B101. In: Bajpal R, Felt DR, Nestler CC, Wani A, Spain JC, et al. (Eds) Federal Integrated Biotreatment Research Consn <7zumC(FIBRC