

Use of Transcutaneous Application of CO₂ in Diabetic Foot Pathology

Igor Frangez^{1*}, Jure Colnaric¹, Danijela Truden²

¹Department of Traumatology, University Medical Centre Ljubljana, Zaloška Cesta, Ljubljana, Slovenia

²Center for Peripheral Vascular Rehabilitation, Kopraska Ulica, Slovenia

*Corresponding Author: Igor Frangez, Department of Traumatology, University Medical Centre Ljubljana, Zaloška Cesta, Ljubljana, Slovenia, Tel: +386 1 522 3255; E-mail: ifrangez@gmail.com

Rec Date: Mar 27, 2017; Acc Date: Apr 12, 2017; Pub Date: April 19, 2017

Copyright: © 2016 Frangez I, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

In neuropathic foot ulcers, the most prominent finding is the loss of peripheral sensation and is typically seen in diabetic patients. In addition, vasculopathy may lead to foot ulcerations in diabetic patients. CO₂ therapy was found to improve chronic wound healing in patients with vascular impairment. It refers to the transcutaneous and subcutaneous application of CO₂ as well as CO₂ water baths for therapeutic purpose. In the method used, gaseous CO₂ is applied transcutaneously using the PVR system®. CO₂ is applied by means of a single-use, low-density polyethylene bag which is wrapped around the leg being treated and secured with an elastic strap. The advantages of this method, compared to injecting CO₂ into subcutaneous tissue, are non-invasiveness, the absence of pain and protection against infection. Compared to CO₂ balneotherapy this approach enables the use of higher CO₂ concentrations, application to chronic wound patients and, with appropriate precautionary measures, prevents the increase of CO₂ in the surrounding air. Finzgar et al. observed that the transcutaneous application of gaseous CO₂ caused a significant increase in the Laser Doppler (LD) flux in cutaneous microcirculation in vivo in humans. The favourable clinical and microcirculatory effects of gaseous CO₂ have further been observed in studies of patients with intermittent claudication as well as patients with primary and secondary Raynaud's phenomenon. The reviewed studies suggest that the increased delivery of CO₂ to the ulcerated area will cause vasodilation and an increase in blood flow. The improved angiogenesis and oxygenation will result in healing of the chronic wound. This principle may be applied in the treatment of diabetic foot ulceration. Moreover, the effect on blood flow may also be important in preventive and curative treatment of patients with impaired mobility due to organic or functional causes. Further work is needed for the development of therapeutic strategies to optimize CO₂ use in diabetic foot patients.

Keywords Carbon dioxide therapy; Microcirculation; Contralateral limb; Vasculopathy; Polyneuropathy

flow in feet to a much higher extent than plain water and it improved the limb salvage rate in critical limb ischemia patients without the option of revascularization. These results showed the potential role of topical CO₂ in effective adjunctive treatment to prevent diabetic ulcer

Carbon dioxide therapy refers to the transcutaneous and subcutaneous application of CO₂ as well as CO₂ water baths for therapeutic purposes. CO₂ rich water bathing has been used since 1930 and was found to improve chronic wound healing in patients with vascular impairment. With neuropathic foot ulcers, the most prominent finding is the loss of peripheral sensation and is typically seen in diabetic patients. Repeated stress and lack of sensation lead to trauma, breakdown of overlying tissue and eventual ulceration. At pressure points, vascular impairment may additionally lead to ulceration.

One major goal of treatment is to improve microcirculation and thereby oxygen supply and the transport of metabolic pathway end-products. The effects of CO₂ therapies on skin microcirculation have been studied in animal models. Duling BR [1] observed increased microvascular diameter and increased perivascular pO₂ at sites of exposure to a CO₂ aqueous solution. The increased diameter was due to the vasodilatory effect of topical CO₂, the phenomenon also known as active hyperaemia. The increased perivascular pO₂ was due to the effect of CO₂ on the oxyhemoglobin dissociation curve. Irie et al. [2] demonstrated that CO₂ immersion induced the production of plasma vascular endothelial growth factor (VEGF), resulting in no-dependent angiogenesis associated with the mobilization of endothelial progenitor cells. Hayashi et al. [3] showed that CO₂ immersion increased blood

LH, *pG l`#0 ð F:c:0€ Q • V

treated using only standard methods. In the group that underwent subcutaneous treatment with CO₂, a significant increase in tissue oxygenation values were observed. They showed progress in healing and a decrease in size of the injured area.

Finzgar et al. [5] conducted a study of the effect of the transcutaneous application of gaseous CO₂ on cutaneous blood flow in vivo in humans. The Laser Doppler (LD) flux in cutaneous microcirculation was measured simultaneously in a group of 33 healthy men during rest and a 35-minute CO₂ therapy. One lower limb of each subject was exposed to gaseous CO₂. The contralateral limb was the control, being exposed to air. The CO₂ therapy caused a statistically significant increase in the LD flux of the studied extremity, whereas in the LD flux of the control extremity was not statistically significant. Aside from a minor decrease in heart rate, no systemic effects were found. The LD flux change is most likely an indirect sign of the successful diffusion of CO₂ molecules through the skin into microcirculation and a direct indicator of the vasodilatory effect of

t

2

a