



White and Brown Rice are Equally Satiating and More Satiating than Glucose Beverage

Xiaochun Snow Wang¹, Mollie O'Neill¹, William Thomas² and Joanne Slavin^{1*}

¹Department of Food Science and Nutrition, University of Minnesota, 1334 Eckles Avenue, Saint Paul, Minnesota 55108, USA

²Division of Biostatistics, School of Public Health, University of Minnesota, 420 Delaware St SE, Minneapolis, MN 55414, USA

Abstract

Liquid meals may evoke weaker appetite signals than calorie-matched solid foods. The objective of this study was to compare the satiety response of white rice and brown rice (whole grain). Additionally, we measured the satiety response of white rice and brown rice compared to a calorie-matched glucose beverage. Each participant (n=20) completed three conditions, presented in random order. On three mornings, separated by at least one week, fasted subjects consumed either 400 kcalories of white rice, brown rice or glucose beverage for breakfast. Visual analogue scales (VAS) were used to assess hunger, satiety, fullness, and prospective food intake at baseline, 15, 30, 45, 60, 90, 120, 180, 240 minutes after breakfast. Gastrointestinal tolerance was assessed at 180 minutes after breakfast and over the 24 hours following each visit using questionnaires. *Ad libitum* lunch was provided at 240 minutes, measured by calorie intake. 24-hour food intake was also recorded by food diary. Satiety differed significantly among treatments, with increased satisfaction and fullness seen with both white rice and brown rice compared to glucose beverage. *Ad libitum* lunch food intake and 24-hour food intake did not differ significantly among treatments. Both white rice and brown rice improve satiety and decrease feelings of hunger more than calorie matched glucose beverage. Enhanced satiety did not translate into reduced food intake at lunch, supporting that many factors may override physiological hunger.

Keywords:

Introduction

1. Liquid meals may evoke weaker appetite signals than calorie-matched solid foods. The objective of this study was to compare the satiety response of white rice and brown rice (whole grain). Additionally, we measured the satiety response of white rice and brown rice compared to a calorie-matched glucose beverage. Each participant (n=20) completed three conditions, presented in random order. On three mornings, separated by at least one week, fasted subjects consumed either 400 kcalories of white rice, brown rice or glucose beverage for breakfast. Visual analogue scales (VAS) were used to assess hunger, satiety, fullness, and prospective food intake at baseline, 15, 30, 45, 60, 90, 120, 180, 240 minutes after breakfast. Gastrointestinal tolerance was assessed at 180 minutes after breakfast and over the 24 hours following each visit using questionnaires. *Ad libitum* lunch was provided at 240 minutes, measured by calorie intake. 24-hour food intake was also recorded by food diary. Satiety differed significantly among treatments, with increased satisfaction and fullness seen with both white rice and brown rice compared to glucose beverage. *Ad libitum* lunch food intake and 24-hour food intake did not differ significantly among treatments. Both white rice and brown rice improve satiety and decrease feelings of hunger more than calorie matched glucose beverage. Enhanced satiety did not translate into reduced food intake at lunch, supporting that many factors may override physiological hunger.

Methods

Subjects

(10 a 10) a 18 36
18 a 27,
E a:
(a a
(<2) ; a a a 5
(a a) ; a
6 ; a
(30 a)
a a ; a a a

*Corresponding author: Joanne Slavin, Department of Food Science and Nutrition, University of Minnesota, 1334 Eckles Avenue, St. Paul, MN 55108, USA, Tel: 612-624-7234; Fax: 612-625-5272; E-mail: jslavin@umn.edu

Received October 21, 2013; Accepted December 12, 2013; Published December 14, 2013

Citation: Wang XS, Neill MO, Thomas W, Slavin J (2013) White and Brown Rice are Equally Satiating and More Satiating than Glucose Beverage. J Obes Weight Loss Ther 3: 201. doi:10.4172/2165-7904.1000201

Copyright: © 2013 Wang XS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

a ; a
a a a a
a (F 1-4).
a a .

Palatability of treatment

a a a a a a a a
a a a . a a a
a a a, a .

Palatability of *ad libitum* lunch

a a a a a , a
a a a, , a a a a . a a a
a a a a a .

Ad Libitum lunch and 24-hour energy intake

a , a a a
a (=0.34) a 24- a (=0.92) a
a (a 1 a F 5).

Gastrointestinal tolerance

re a a
a a . A
a Gi a (a 1). Gi a

a a a a 24 (=0.026).
a a a H₂ a 24- Gi- a
a .

Discussion

a a a , a
a a a a a , a
a a a
a

a a a - a 18 . G
19 a a a a a a
a a a a a a a . a . 6
a a a a a a a

a 2 a a a a 2 a i a
a a a a a a a a a
a a a a a a a a a
a 14 . a a a (. .)
a a a a a a a a
A A a a a a
a 24- a a a a a a a a
a a a a a a a a a
a a a a a a a a a
a a a a a a a a a
E (887 a), (900 a) a a a
(959 a) , a a a a a a a a a
a a a a a a a a a a a
17 . A 28- a a a a a a a
a a a a a a a a a a
a B a a a a a a a a a
a a a a a a a a a a
a a a a a a a a a a

3. Itani T, Tamaki M, Arai E, Horino T (2002) Distribution of amylase, nitrogen, and minerals in rice kernels with various characters. *J Agric Food Chem* 50: 5326-5332.
4. Stull AJ, Apolzan JW, Thalacker-Mercer AE, Iglay HB, Campbell WW (2008) Liquid and solid meal replacement products differentially affect postprandial appetite and food intake in older adults. *J Am Diet Assoc* 108: 1226-1230.
5. Flood-Obbagy JE, Rolls BJ (2009) The effect of fruit in different forms on energy intake and satiety at a meal. *Appetite* 52: 416-422.
6. Willis HJ, Thomas W, Willis DJ, Slavin JL (2011) Feasibility of measuring gastric