mono lament examination, it is possible that these new tingling sensations were initially interpreted as pain. Unfortunately we did not attempt to distinguish di erences between tingling and pain. By the 12th week, NRS pain levels were ve points (6.0 versus 1.0) lower when compared to the values reported during the rst week of the intervention (Figure 1) and correspond with the decreases observed from the BPI-sf pain severity index (Table 1). is is considered clinically signi cant given the change of pain severity classi cation from moderate (4-6) to mild (1-3) [6,13].

Although we do not identify a speci c mechanism for the change in the sensory aspect of pain, painful symptoms related to neuropathy in diabetics have been strongly associated with poor blood glucose control [14]. Our patient noted signi cant improvements in her fasting glucose and HbA1c or glycated hemoglobin (Table 1). Hemoglobin A1c levels between 5.7% and 6.4% indicate increased risk of diabetes whereas, levels between 4% and 5.6% are considered normal. Our patients, Hemoglobin A1c level decreased from a baseline measure of 6.1 to 5.1 at the conclusion of the intervention. e decrease our patient observed in her HbA1c support the ndings of Lee et al. [15] who also report signi cant decreases in HbA1c in the elderly with DNP following WBV training.

Our results indicate that consistent use of WBV is associated with an increase in walking function, as de ned by walking speed. Our patients' baseline walking speed of 1.068 m/s classi ed her at a higher risk for all cause mortality. Women with walking speeds <1.09 m/s are associated with a higher risk of death [12]. Post-intervention, her walking speed increased to 1.214 m/s, a 13.7% improvement thus reducing her risk of death. In healthy older adults, a change in walking speed of 0.05 m/s is considered clinically meaningful [16], our patient observed an increase in walking speed of 0.146 m/s. is improvement is consistent with changes observed in elderly adults [17] and individuals with Parkinson's disease [18] receiving WBV therapy. e application of WBV has been shown to increase muscular activation [19], strength and power [20,21], peripheral blood ow circulation [22-24], acutely decreases arterial sti ness in healthy young men (age 26.6 ± 1.9 years) [25], and acutely improve skin blood ow when directly applied to the forearm in diabetics [24] Additionally, WBV has been shown to increase blood ow and activate muscle activity in patients with Friedreich's ataxia and neurodegenerative disease [26] that causes damage to the nervous system. With increased muscle activity, blood ow rate to the active muscles will increase by as much as 20 times the resting rate. We attribute the improved walking function directly to the signi cant decrease in pain and edema, potentially a result of improved blood circulation in the lower legs resulting from the WBV therapy.

At the completion of the WBV intervention, our participant reported that as the pain in her feet decreased, her physical activity levels increased and she eventually joined a tness center. is in turn, lead to other signi cant health bene ts including a decrease in fasting glucose (121 to 106), hypertriglyceridemia (drop of 287 points), and a decrease in peripheral edema. Moreover, her pain and hypertension symptoms improved to the extent that her Lyrica and Benazepril HCl were stopped, and her Metformin was reduced by 50%. Additionally, our patient observed a weight loss (15.91 kg) and decrease in percent body fat of 36.5%. We have shown experimentally that WBV has bene cial e ects related to weight gain prevention in female rats [27].

C 1 1

is case study provides rst evidence that WBV could be an e ective and relatively inexpensive therapeutic modality compared to prescription drugs to treat patients with painful diabetic neuropathy. Of course, appropriately powered studies are needed to fully elucidate its therapeutic potential. However, the magnitude and broad spectrum of improvements provides a basis to expeditiously embark on such an investigation.

Acknowledgment

John C. Erkkila, M.D. Endowment For Health and Human Performance.

References

1. NIH N.I.O.H. (2009) Diabetic neuropathies: The nerve damage of diabetes. NIH Publication No. 09-3185.
2. Rittweger J (2010) Vibration as an exercise modality: how it may work, and what its potential might be. Eur J Appl Physiol 108: 877-904.
3. Alentorn-Geli E, Padilla J, Moras G, Lázaro Haro C, Fernández-Solà J (2008) Six weeks of whole-body vibration exercise improves pain and fatigue in
4. Iwamoto J, Takeda T, Sato Y, Uzawa M (2005) Effect of whole-body vibration exercise on lumbar bone mineral density, bone turnover, and chronic back pain in post-menopausal osteoporotic women treated with alendronate. Aging Clin Exp Res 17: 157-163.
5. Dworkin RH, Turk DC, Wyrwich KW, Beaton D, Cleeland CS, et al. (2008) Interpreting the clinical importance of treatment outcomes in chronic pain clinical trials: IMMPACT recommendations. J Pain 9: 105-121.
6. Deborah L. Hoffmana, Alesia Sadosky B, Ellen M. Dukesb, Jose Alvirb (2010) How do changes in pain severity levels correspond to changes in health status and function in patients with painful diabetic peripheral neuropathy? Pain 149: 194-201.
7. Pain and its treatment in outpatients with metastatic cancer. N Engl J Med 330: 592-596.
8. Cleeland CS, Ryan KM (1994) Pain assessment: global use of the Brief Pain Inventory. Ann Acad Med Singapore 23: 129-138.
9. Jensen MP, Chodroff MJ, Dworkin RH (2007) The impact of neuropathic pain on health-related quality of life: review and implications. Neurology 68: 1178-1182.
10. Cooper R, Kuh D, Hardy R; Mortality Review Group; FALCon and HALCyon Study Teams (2010) Objectively measured physical capability levels and mortality: systematic review and meta-analysis. BMJ 341: c4467.
11. Dumurgier J, Elbaz A, Ducimetière P, Tavernier B, Alpérovitch A, et al. (2009) Slow walking speed and cardiovascular death in well functioning older adults: prospective cohort study. BMJ 339: b4460.
12. Elbaz A, Sabia S, Brunner E, Shipley M, Marmot M, et al. (2013) Association of walking speed in late midlife with mortality: results from the Whitehall II cohort study. Age (Dordr) 35: 943-952.
13. Diane C. Zelman, Ellen Dukes, Nancy Brandenburg, Alan Bostrom, Mugdha