Open Access

The Mast I 一鱅粔aQ竻 7 碆脟蕒湣箲虱冠! 曷箲擭荋

A Drissi Maniani Abdelilah^{1*}, S.habibchorfa¹, S Essetti Latib Rachida¹, OmorYoussef¹, Karima el Matara², Merbou Sahar S² and Jellal H. Jerguigue²

Department of Radiology, Faculty of Medicine and Pharmacy of Rabat, Oncology National Institute, IbnSina University Hospital, Rabat, Morocco

Abstract

The mastitis carcinomatosa or infammatory breast is an aggressive form of T usual diferent histologi mainly based on clinical and radiological criteria, MRI is more reliable in assessing tumor size than clinical examination, The most common presentation is a poorly limited heterogeneous infltrating mass associated with thickening. The 5-year survival rate range 19-59% depends on the tumor stage. The treatment of infammatory breast cancer is multidisciplinary, combining neoadjuvant chemotherapy followed by loco regional treatment.

Solutions of the solution of t

 C_{1} ,

and the first of the second se enting to colorate with the colorate beauties.

- the colorate term of the first colorate beauties.

- the colorate term of the first colorate beauties.

- the colorate term of the first colorate beauties.

*Corresponding author: A Drissi Maniani Abdelilah, Department of Radiology, Faculty of Medicine and Pharmacy of Rabat, Oncology National Institute, IbnSina University Hospital, Rabat, Morocco, Tel: 212660323703; E-mail: abdel2020abrar@

Received: 18-Dec-2021, Manuscript No. roa-21-50169; Editor assigned: 20-Dec-2021, PreQC No. roa-21-50169(PQ); Reviewed: 12-Jan-2022, QC No. roa-21-50169; Revised: 17-Jan-2022, Manuscript No. roa-21-50169(R); Published: 24-Jan-2022, DOI: 10.4172/2167-7964.1000360

Citation: Abdelilah ADM, Habibchorfa S, Rachida SEL, Youssef O, Matara K, et al. (2022) The Mastitis Carcinomatosa, a Case Report. OMICS J Radiol 11: 360.

Copyright: © 2022 Abdelilah ADM. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

²Department of Gynecology-obstetrics, Oncology and High-risk pregnancy Souissi Rabat

I whaten a should be took at the following to be such to be a should be a shou

المرافعة ال مرافعة المرفعة مرافعة المرفعة المرفعة

الماري المار الماري الماري