



## A Note on Fetal Macrosomia

**Nidhi Shah\***

*American Board of Pediatrics, Texas Tech University Health Sciences Center at Amarillo, USA*

---

**\*Corresponding author:**

**A**

I would like to thank my Professor for his support and encouragement.

**C I**

The authors declare that they are no conflict of interest.

#### References

1. Shah Z, Chawla A, Patkar D, Pungaonkar S (2003) "MRI in kernicterus". *Australas Radiol* 47: 55-57.
2. Gómez M, Bielza C, Pozo FD, Rios Insua S (2007) "A graphical decision-theoretic model for neonatal jaundice". *Med Decis Making* 27: 250-265.
3. Gilmour, Susan M (2004) "Prolonged neonatal jaundice: When to worry and what to do". *Paediatr Child Health* 9: 700-704.
4. Duryea EL, Hawkins JS, McIntire DD, Casey BM, Leveno KJ (2014) A revised birth weight reference for the United States. *Obstet Gynecol* 124: 16-22.
5. Nesbitt TS, Gilbert WM, Herrchen B (1998) Shoulder dystocia and associated risk factors with macrosomic infants born in California. *Am J Obstet Gynecol* 179: 476-480.
6. Boulet SL, Alexander GR, Saliyu HM, Pass M (2003) Macrosomic births in the United States: determinants, outcomes, and proposed grades of risk. *Am J Obstet Gynecol* 188: 1372-1378.
7. Zhang X, Decker A, Platt RW, Kramer MS (2008) How big is too big? The perinatal consequences of fetal macrosomia. *Am J Obstet Gynecol* 198: 517.
8. Colletti JE, Kothari, S, Kothori S, Jackson DM, Kilgore KP, et al. (2007) "An emergency medicine approach to neonatal hyperbilirubinemia". *Emerg Med Clin North Am* 25: 1117-35.
- 9.