

Adverse Effects of Acid Rain on Living Things

Mohammad Shain*

Department of Natural Resources and Environment, Arabian Gulf University, Bahrain

Abstract

Rain or any other type of precipitation that is unusually acidic is known as acid rain because it contains more hydrogen ions (low pH) than normal. The majority of water, including drinking water, has a neutral pH between 6.5 and 8.5; however, acid rain typically has a pH level between 4 and 5. The pH of the acid rain decreases with increasing acidity. Infrastructure, aquatic animals, and plants can all be damaged by acid rain. Sulfur dioxide and nitrogen oxide emissions, which react with the water molecules in the atmosphere to produce acids, are what lead to acid rain.

Keywords: Acid rain; Air pollution

Introduction

It has been demonstrated that acid rain has negative effects on forests, freshwaters, soil, microbes, insects, and aquatic life. Consistent acid rain weakens the durability of tree bark in ecosystems, making plants more vulnerable to drought, heat/cold, and pest infestation. Corrosive downpour is likewise a threat for detriming soil synthesis by stripping it of supplements, for example, calcium and magnesium which assume a part in plant development and keeping up with sound soil. In addition to having an effect on human health, acid rain also causes paint to peel, steel structures like bridges to corrode, and stone buildings and statues to weather.

A few legislatures, remembering those for Europe and North America, have put forth attempts since the 1970s to decrease the arrival of sulfur dioxide and nitrogen oxide into the environment through air contamination guidelines. These endeavors have had positive outcomes because of the far and wide exploration on corrosive downpour beginning during the 1960s and the plugged data on its hurtful impacts. Sulfur dioxide and nitrogen oxides can also be produced naturally by lightning, and volcanic eruptions are the primary sources of sulfur and nitrogen compounds that cause acid rain. Although acid rain has a much smaller global impact on oceans, it has a greater impact in the shallower waters of coastal waters. Acid rain can cause the pH of the ocean to decrease, a process known as ocean acidification, making it more difficult for various coastal species to construct the exoskeletons they require to survive. Coral's limestone skeleton is particularly sensitive to pH decreases because calcium carbonate, a core component of the limestone skeleton, dissolves in acidic (low pH) solutions. These coastal species link together as part of the ocean's food chain; without them, more marine life will die [1-4].

Discussion

Excess atmospheric nitrogen contributes to both acidification and phytoplankton and other marine plant growth, which may increase the frequency of harmful algal blooms and eutrophication (the formation of oxygen-depleted "dead zones" in some parts of the ocean).

Fish and other aquatic animals may suffer harm as a result of the acid rain's lower pH and higher aluminum concentrations in surface water. The majority of fish eggs will not hatch at a pH below 5, and lower pH can kill adult fish. The diversity of lakes and rivers decreases as they become more acidic. In some lakes, streams, and creeks in geographically sensitive areas, such as the Adirondack Mountains in the United States, acid rain has eliminated insect life and some fish species, including brook trout. However, the extent to which acid

rain directly or indirectly contributes to lake and river acidity (i.e., depending on characteristics of the surrounding watershed) is variable.

*Corresponding author: Mohammad Shain, Department of Natural Resources and Environment, Arabian Gulf University, Bahrain, E-mail: shainmd@edu.bh

Received: 05-Dec-2022, Manuscript No. EPCC-22-84067; **Editor assigned:** 07-Dec-2022, PreQC No. EPCC-22-84067 (PQ); **Reviewed:** 21-Dec-2022, QC No. EPCC-22-84067; **Revised:** 23-Dec-2022, Manuscript No. EPCC-22-84067 (R); **Published:** 30-Dec-2022, DOI: 10.4172/2573-458X.1000316

Citation: Shain M (2022) Adverse effects of Acid Rain on Living Things. Environ Pollut Climate Change 6: 316.

Copyright: © 2022 Shain M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

of the plant's existence. Adverse effects, such as the acid's effects on soil (see above) or a high concentration of gaseous acid rain precursors may have an indirect connection to acid rain. Forests at high altitudes are particularly vulnerable because they are frequently surrounded by fog and clouds that are more acidic than rain.

The application of lime and fertilizers to replace nutrients minimizes the impact on food crops, which can also be damaged by acid rain. In developed regions, limestone may likewise be added to build the capacity of the dirt to keep the pH stable, however this strategy is generally unusable on account of wild terrains. Red spruce trees become less tolerant of cold and are more likely to suffer injury or even die during the winter if calcium is leached from their needles. Western nations have been coming to an agreement on international treaties regarding the long-distance transport of atmospheric pollutants for some time. Beginning in 1979, nations from Europe gathered to ratify the UNECE Convention's general principles. The goal was to stop Long-Range Transboundary Air Pollution [9-10].

Conclusions

The Helsinki Protocol on the Reduction of Sulphur Emissions under the Convention on Long-Range Trans boundary Air Pollution was a way to make the convention's results better. All effects of the deal have proactively worked out as expected, as confirmed by a rough 40 percent drop in particulate matter in North America. Additional international commitments to stop the spread of particulate matter have been inspired by the effectiveness of the Convention in combating acid rain. In 1991, the Air Quality Agreement was signed by the US and Canada. The treaties were signed by Canada and most European nations. Movement of the Long-Reach Trans boundary Air Contamination Show stayed lethargic after 1999, when 27 nations met to additionally decrease the impacts of corrosive rain. In 2000,

unfamiliar collaboration to forestall corrosive downpour was started in Asia interestingly. Ten diplomats from various nations on the continent met to talk about ways to stop acid rain.

References

1. Laden F, Schwartz J, Speizer F, Dockery D (2006) Reduction of Long-Range Transboundary Air Pollution in Asia. *Environmental Health Perspectives* 114: 1033-1040.