

compression stockings to help prevent clots from forming.

Conclusion

In summary, the symptoms of VTE can vary depending on the location and severity of the clot, and treatment typically involves anticoagulation therapy and/or mechanical interventions to prevent the clot from growing and to reduce the risk of future clots. Prevention is also important for individuals who are at high risk of VTE.

Acknowledgement

None

Conflict of Interest

None

1. Morris TA (2011) Natural history of venous thromboembolism. *Circulation* 124: 869-884.
2. Geerts WH, Bergqvist D, Pineo GF, Heit JA, Samama CM, et al. (2008) Prevention of venous thromboembolism: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest* 133: 381S-453S.
3. Essien EO, Rali P, Mathai SC (2004) Pulmonary embolism. *Med Clin North Am* 78: 594-564.
4. Heit JA, Cohen AT, Anderson FA Jr (2005) Estimated annual number of incident

and recurrent, non-fatal and fatal venous thromboembolism (VTE) events in the US. *Blood* 106: 910.

5. Becattini C, Agnelli G (2016) Risk stratification and management of acute pulmonary embolism. *Hematology. Am Soc Hematol Educ Program* 2016: 404-412.
6. Lutsey PL, Zakai NA (2023) Epidemiology and prevention of venous thromboembolism. *Essentials in Hematology, Cardiology* 6(2): 248-261.