

Keywords: Diabetes mellitus; Retina; Hypertension; High cholesterol; Diabetic retinopathy

I

Diabetic retinopathy is a common and potentially sight-threatening complication of diabetes mellitus. It affects the retina, the light-sensitive tissue at the back of the eye, and is a leading cause of blindness among adults worldwide. As the prevalence of diabetes continues to rise, diabetic retinopathy becomes an increasingly significant public health concern. This article provides a short overview of diabetic retinopathy, its causes, symptoms, risk factors, and the importance of early detection and management [1].

Ca

Diabetic retinopathy is primarily caused by long-term uncontrolled high blood sugar levels. Elevated glucose levels in the blood can damage the small blood vessels that nourish the retina, leading to leakage, swelling, and eventually, the growth of abnormal blood vessels. These abnormal vessels can leak blood into the retina, causing further damage and vision loss.

S

In its early stages, diabetic retinopathy may not cause noticeable symptoms. [2] Blurred or distorted vision

- Fluctuating vision
- Floaters (small, dark spots or strings in the field of vision)
- Impaired color perception
- Dark or empty areas in the visual field

Risk factors

Several factors increase the risk of developing diabetic retinopathy:

- Poorly controlled diabetes: People with consistently high blood sugar levels are at greater risk.
- Duration of diabetes: The longer a person has diabetes, the higher the risk of diabetic retinopathy.
- High blood pressure: Hypertension can exacerbate retinal damage.
- High cholesterol levels: Elevated cholesterol can contribute

to vascular problems in the eye.

- Pregnancy: Pregnant women with diabetes may experience worsened retinopathy [3].
- Ethnicity: Certain ethnic groups, such as Hispanics, African-Americans, and Native Americans, are more susceptible to diabetic retinopathy.

Prevention

- Early detection and proactive management are crucial in preventing vision loss due to diabetic retinopathy.
- Regular eye exams: Annual comprehensive eye examinations by an ophthalmologist are essential for early detection and intervention.
- Blood sugar control: Maintaining tight control of blood glucose levels can slow the progression of diabetic retinopathy.
- Blood pressure management: Keeping blood pressure within a healthy range is vital for preserving eye health [4].
- Cholesterol control: Monitoring and managing cholesterol levels can reduce the risk of vascular complications.
-

and identify signs of retinopathy.

O a a (OCT): OCT is a non-invasive imaging technique that provides high-resolution cross-sectional images of the retina. It helps detect retinal swelling, fluid accumulation, and other abnormalities [5].

F a a : In this procedure, a fluorescent dye is injected into a vein in the arm, and images are taken as the dye circulates through the blood vessels in the retina. This helps identify any leakage or abnormal blood vessel growth.

M a a a a

B a : Maintaining stable blood sugar levels is crucial in preventing or slowing the progression of diabetic retinopathy. This involves adherence to a well-balanced diet, regular exercise, and appropriate diabetes medications.

B a a : Controlling hypertension is essential as high blood pressure can exacerbate retinal damage.

C a a : Monitoring and managing cholesterol levels help reduce the risk of vascular complications in the eye.

I a a : In advanced stages of diabetic retinopathy, intravitreal injections of medications, such as anti-VEGF agents or steroids, may be administered to reduce swelling and control abnormal blood vessel growth [6].

L a a a : Laser treatment is often used to seal leaking blood vessels and prevent further damage. It can also be employed to shrink abnormal blood vessels.

V : In severe cases with significant bleeding or tractional retinal detachment, vitrectomy surgery may be necessary to remove the vitreous gel and repair the retina.

R a w - : Patients with diabetic retinopathy require regular follow-up visits with their eye care specialists to monitor the condition and adjust treatment as needed.

R

Diabetic retinopathy is a serious eye condition resulting from long-term uncontrolled diabetes. It affects the blood vessels in the retina, leading to vision impairment and, if left untreated, eventual blindness.

The condition develops gradually and often remains asymptomatic in its early stages, making regular eye check-ups crucial for early detection [7].

As diabetes prevalence continues to rise globally, diabetic retinopathy is becoming a significant public health concern. The risk factors for developing the condition include poorly controlled blood sugar levels, high blood pressure, and prolonged diabetes duration. Individuals from certain ethnic backgrounds also face a higher risk.

Prompt intervention is essential in managing diabetic retinopathy. Treatment methods include laser photocoagulation to seal leaking blood vessels and intravitreal injections to reduce swelling. Maintaining stable blood sugar levels, managing blood pressure, and adopting a healthy lifestyle can slow the progression of the disease.

Diabetic retinopathy highlights the importance of comprehensive diabetic care and regular eye examinations. Early diagnosis and proactive management are crucial for preserving vision and improving the quality of life for individuals with diabetes. Public awareness and timely intervention are key to mitigating the impact of this vision-

threatening complication [8].

D

The rising prevalence of diabetes worldwide has resulted in an increased number of individuals at risk of developing diabetic retinopathy. The disease progresses gradually, and its early stages may be asymptomatic, which emphasizes the need for regular eye examinations among diabetic patients. Early detection plays a critical role in preventing irreversible damage to the retina and preserving vision [9].

The primary cause of diabetic retinopathy is prolonged exposure to elevated blood sugar levels, which damages the delicate blood vessels that nourish the retina. This damage results in leakage, swelling, and the formation of abnormal blood vessels, leading to impaired vision. While blood sugar control remains the cornerstone of diabetes management, it is essential to recognize other risk factors such as high blood pressure and cholesterol levels, which can exacerbate retinal damage.

Managing diabetic retinopathy requires a multidisciplinary approach. Ophthalmologists, endocrinologists, and primary care physicians must work together to provide comprehensive care for patients with diabetes. Early intervention through laser photocoagulation or intravitreal injections can help control the progression of the disease and prevent further vision loss. Additionally, maintaining a healthy lifestyle, including a balanced diet, regular exercise, and abstaining from smoking, can significantly improve outcomes.

Public awareness campaigns are crucial in ensuring that individuals with diabetes understand the potential risks of diabetic retinopathy and the importance of regular eye examinations. [10] Timely intervention can prevent severe complications, reduce healthcare costs, and enhance the overall quality of life for affected individuals.

C

hospitalized children in Abadan, Iran. Gastroenterol Hepatol Bed Bench 7: 218.

2.