

## Cadmium Induced Toxicity of Liver

Emma Quile\*

Department of toxicology, Horbachevsky Ternopil State Medical University, Ukraine

### Abstract


Cadmium is a harmful component to what man can be uncovered working or in the climate. Album's most striking toxicological property is its particularly lengthy half-life in the human body. When retained, Album collects in the human body, especially in the liver. The cell activities of Cd are widely archived, however the sub-atomic instruments basic

Toxicol 6: 213.

**Copyright:** © 2023 Quile E. This is an open-access article distributed under the

$\gamma\theta$ Hepatic Disc content expanded step by step with age without a sharp fall, as did kidney Cd substance. hward breath of cadmium oxide vapor sprayers can cause intense pneumonitis and aspiratory edema. Nonetheless, in persistent word related or ecological openness, Cd can

## Conclusion



## References

1. Toxicogenomics-based discrimination of toxic mechanism in HepG2 human hepatoma cells. *Toxicol Sci* 58: 399-415.
2. Optimization of microarray data analysis in the cladoceran *Daphnia magna*.
3. Yeast microarrays for genome wide parallel genetic and gene expression analysis.
4. DNA microarrays and toxico-genomics: applications for ecotoxicology? *Biotechnol Adv* 20: 391-419.
5. Renn SC, Aubin-Horth N, Hofmann HA. 2004. Biologically meaningful microarray. *BMC Genom* 5: 42.
6. Schena M, Shalon D, Heller R, Chai A, Brown PO et al. (1996) Parallel human genome analysis: microarray-based expression monitoring of 1000 genes. *Proc Natl Acad Sci USA* 93: 10614-10619.
7. Overview on microarray data analysis in toxicology assessment. *Environ Health Perspect* 112: 460-464.
8. Genomic microarrays in toxicology. *Environ Health Perspect* 111: 825-835.
9. Standards for microarray data analysis.
10. Monitoring the effects of salinity stresses using a full-length cDNA microarray.