



Neonatal jaundice is a common condition in newborn babies, characterized by yellowing of the skin and eyes. It is caused by a buildup of bilirubin in the blood. The most common cause is an imbalance between bilirubin production and excretion. In newborns, the liver is still developing and may not be able to process bilirubin efficiently. Other factors that can contribute to jaundice include breastfeeding, dehydration, and certain medications. Jaundice is usually diagnosed by a physical examination and confirmed by a blood test. Treatment typically involves phototherapy, which uses blue light to break down bilirubin in the skin. In severe cases, an exchange transfusion may be necessary. Most cases of jaundice resolve within a few days, but it is important to monitor the baby's bilirubin levels closely to prevent complications.

Neonatal jaundice is a common condition in newborn babies, characterized by yellowing of the skin and eyes. It is caused by a buildup of bilirubin in the blood. The most common cause is an imbalance between bilirubin production and excretion. In newborns, the liver is still developing and may not be able to process bilirubin efficiently. Other factors that can contribute to jaundice include breastfeeding, dehydration, and certain medications. Jaundice is usually diagnosed by a physical examination and confirmed by a blood test. Treatment typically involves phototherapy, which uses blue light to break down bilirubin in the skin. In severe cases, an exchange transfusion may be necessary. Most cases of jaundice resolve within a few days, but it is important to monitor the baby's bilirubin levels closely to prevent complications.