Keywords:

Introduction

Advances and research in oral pathology

Digital pathology:

Targeted therapies:

Regenerative medicine:

Conclusion

References

- Tun KM, Imwong M, Lwin KM, Win AA, Hlaing TM, et al. (2015) Spread of artemisinin-resistantPlasmodium falciparumin Myanmar: a cross-sectional survey of the K13 molecular marker. THE LANCET Infectious Diseases 15: 415-421.
- Akoria OA, Arhuidese IJ (2014) Progress toward elimination of malaria in Nigeria: Uptake of Artemisinin-based combination therapies for the treatment of malaria in households in Benin City. Annals of African medicine 13: 104-113
- Baragana B, Hallyburton I, Lee MCS, Norcross NR, Grimaldi R, et al. (2015) A novel multiple-stage antimalarial agent that inhibits protein synthesis. Nature 522: 315-320.
- Exavery A, Mbaruku G, Mbuyita S, Makemba A, Kinyonge IP, et al. (2014) Factors afecting uptake of optimal doses of sulphadoxine-pyrimethamine for intermittent preventive treatment of malaria in pregnancy in six districts of Tanzania. Malaria Journal 13: 10-1186.
- Simba DO, Kakoko D, Tomson G, Premji Z, Petzold M, et al. (2012) Adherence to artemether/lumefantrine treatment in children under real-life situations in rural Tanzania. Transactions of the Royal Society of Tropical Medicine and Hygiene 106: 3-9.
- Bruxvoort K, Kalolella A, Cairns M, Festo C, Kenani M, et al. (2015) Are Tanzanian patients attending public facilities or private retailers more likely to adhere to artemisinin-based combination therapy? Malaria Journal 14: 1-12.
- Win TZ, Zaw L, Khin W, Khin L, Tin OM, et al. (2012) Adherence to the recommended regimen of artemether-lumefantrine for treatment of uncomplicated falciparum malaria in Myanmar. Myanmar Health Science Research Journal 24: 51-55.
- Zbe OP, Mangham-Jeferies L, Cundill B, Wiseman V, Uzochukwu BS, et al. (2015) Quality of care for the treatment for uncomplicated malaria in South-East Nigeria: how important is socioeconomic status?. International Journal for Equity in Health 14: 19.
- Watsierah CA, Jura WG, Oyugi H, Abong'o B, Ouma C (2010) Factors determining anti-malarial drug use in a peri-urban population from malaria holoendemic region of western Kenya. Malar J 9: 295.
- Das A, Dash A (2007) Evolutionary paradigm of chloroquine-resistant malaria in India. Trends Parasito I23: 132-135.