

Viral Coinfections

Youssef Moutaouakkil ^{1*}, Mohamed Reda Tagajdid ², Mohammed Sbaai ¹, Safae Elkochri ¹, Rachid Abi ¹, Hicham El Annaz ¹ and Saad Mrani ²

¹HSDUWPHQW RI 9LUORJ\ ORKDPHG 9 OLOLWDU\ 7HDFKLQJ +RVSLWDO ORKDPHG 9 8QLYHUVLW\ 5DEDW +XPDQ DQG 0ROHFXODU 9LUORJ\ ²HSDUWPHQW ORKDPHG 9 OLOLWDU\ 7HDFKLQJ +RVSLWDO ORKDPHG

Keywords: Hepatitis C; HIV; Liver; Carcinoma

Coinfection HBV/HCV

The hepatitis C virus belongs to the family Flaviviridae. It is estimated that 130 million people are infected with HCV worldwide and that cases of co-infections HBV / HCV concern 7-20000000 people. These two viruses share the same liver tropism. The case of co-infections is associated with increasing severity of liver disease, faster development to stages of fibrosis, cirrhosis and HCC than in the case of monoinfection whether HBV or HCV [1]. Studies conducted in vitro have shown that HCV played a dominant negative effect which inhibit HBV replication and which is mediated by the HCV Core protein [2,3]. A second study, meanwhile, found the opposite effect, suggesting that HBV inhibit replication of HCV [4]. However, or interactions between HBV and HCV are still very difficult to study and this mainly because of lack of appropriate studies in vitro systems. BELLECABE team was able to demonstrate that these two viruses could replicate in the same cell without bias interact directly. Viral interference would be indirect and mediated by molecular mechanisms that might be responsible for these mutual inhibitions observed [5-8]. HCV is capable of inducing an

*Corresponding author: <R XMV 0RXWDRXDNNLO 'HSDUWPHQW RI 9 OLOLWDU DBKLQJ +R R&DWBG 908QLYHUVLW\ 7DDEDW ORL (PDLO \RXV VHI PE #KRWPDLQ I U

Received: 'HF HPEHU Accepted:)HEUXDU\ Published)HEUXDU\

Citation: 0RXWDRXDNNDQGLG E DDL 0 (O\$IRIFKULHW DO 9LU DO &RLQIHFWRQV \$G2Y, 0RO 'LDJ

Copyright: < 0RXWDRXDNNDQGLG 7KLV LV DQ RSHQ DFFHV XQGHU WKH WHUPV RI WKH &UHDWLYH &RPPRQV \$W XQUHVWXULHF WBGVVWULEXWLQRQ DQG UHSURGXFWLRQ RULJLQDO DXWKRU DQG VRXUFH DUH FUHGLWHG

at key positions Y100C, P120T, sK122R, sD144A. In two studies are reported sK122R mutations and involve genotypes A, substitutions extremely rare in this genotype. The absence of antibody at the time of reactivation was not possible to perform functional analysis of the recognition of HBsAg and anti-HBs antibodies of patients. One can imagine that the appearance of mutation is the result of immune selection pressure on the virus. But that reactivation of HBV was made possible by the absence of anti-HBs antibodies. The reactivation of occult infections in HIV patient's phenomena (+). Another study showed that reactivation of occult infection in two HIV (+) patients was independent of resistance mutations to lamivudine and state of immunosuppression [18].

References

- 3RWWKRII \$ 0DQQV 03 :HGHTPHWPHQ%9 R&9
FRLQIHFWRQ ([SHUW 2SLQ 3KDUPDFRKHU
6KLK &0 6- /R 07L\BRMQD6< /HH <+ 6XSSUHVVL
% YLUJV H[SUHVVLRQ DQG UHSOLFDWL RQ E\ KHS
FHOOV - 9LURO
6FKXWWOHU &* 6FLKIRGLOGHWU . 5HSS 5 * HUOLFK :+
6XSSUHVVLRQ RI KHS DWL WL V % YLUJV H
SURWHLQ - +HSDWRO
3RQWL VVR 30 5XDWVWHRW MFRK ** &KPHHOOR / * DOORULQ H W DO
*DVWURHQWHURORJ\\
.RLNH<DVXGD . <RWVX\IDQDJL + ORUL\D . + LQR .
UHSOLFDWL RQ RI HLWKHU YLUJV LQ GXDO LQIHF
9LURO
%HOOHFDRYHWHQRLUH - *DMHU 0 %UDVV 9 .RXWVRXGDNLV * HW DO
+HSDWLWL V % DQG & YLUJV FRLQIHFWRQ \$ QRYHO PRGHO V\VWHP UHYHDOV WKH DEVHQFH
RI GLUHF YLUJO LQWHUIHUhQFH +HSDWRORJ\\
(\UH 16KLOOLSV 5- %RZGHQ 6 <LS ('HZDU % HW DO
+HSDWLWL V % YLUJV