Open Access Scientific Reports

Research Article Open Acces

Abstract

Keywords: RabGGTase; Docking; Prenyltransferases; OPLS-AA force eld; Pyrimidine; QuickPro; GLIDE

Introduction

Rab proteins are small GTPases involved in the regulation of vesicular tra c [1]. To be functional, they have to undergo a post-translational modi cation in which one or two isoprenoid geranylgeranyl groups are attached to their C termini [2]. Rab geranylgeranyltransferase (RabGGTase or GGTase type II) catalyzes this reaction. RabGGTase belongs to a family of prenyltransferases that includes farnesyltransferase (FTase) and geranylgeranyltransferase type I. ese enzymes are highly conserved in sequence and structure and possess a single active site [3].

Pyrimidines are an important class of heterocyclic compounds, which possess a wide range of biological activities such as anticancer [4,5], antibacterial [6], anti-in ammatory [7], antiviral [8], antitubercular [9], antihypertensive [10] and anticonvulsant [11] activities. e presence of 4-OCH_3 and 2-NO_2 and 3-NO_2 substitutions on pyrimidines shows signi cant increase in antitubercular activity [12]. e bromo substituted, 2-hydroxy phenyl substituted pyrimidines also shows good antihistaminic activity [13].

To know the possibility of binding of RabGGTase protein to pyrimidine analogues is fascinating and molecular docking is frequently used to predict the binding orientation. Docking is used to bring out new ligands for biological targets with a known 3D structure [14]. Docking studies of the compounds were performed using GLIDE (Schrödinger, LLC, Portland) [15,16].

Materials and Methods

Ligand structure preparation

e structures of six pyrimidine analogues, a) 4-(2- uoro-4-methoxyphenyl)-6-(2,3, trimethoxyphenyl) pyrimidin-2-amine, b) 4-(2- uoro-4-methoxyphenyl)-6-(2,4,6-trimethoxyphenyl) pyrimidin-2-amine, c) 4-(2- uoro-4-methoxyphenyl)-6-(2, 4-dimethoxyphenyl) pyrimidin-2-amine, d) 4-(2- uoro-4-methoxyphenyl)-6-(2,4,5-trimethoxyphenyl) pyrimidin-2-amine, e) 4-(2- uoro-4-methoxyphenyl)-6-(3,4,5-trimethoxyphenyl) pyrimidin-2-amine, f) 4-(2- uoro-4-methoxyphenyl)-6-(3,5-dimethoxyphenyl) pyrimidin-2-amine (Figure 1) were constructed using the fragment dictionary of Maestro 7.5 (Schrodinger, LLC) using the Optimized Potentials for Liquid Simulations-All Atom (OPLS-AA) force eld [17] with the steepest descent followed by truncated Newton conjugate gradient protocol. Partial atomic charges were computed using the OPLS-AA force eld. e ligand molecules and their methoxy substitution are shown in table 1.

*Corresponding author: Sajitha Lulu S, Assistant Professor, School of Biosciences and Technology, VIT University, Vellore, Tamil Nadu-632014, India, Tel: +91 9944807641; E-mail: ssajithalulu@vit.ac.in

Received April 29, 2011; Published January 15, 2013

Citation: Ethiraj KR, Vino S, Nawas Khan F, Sajitha Lulu S (2013) Docking Studies of Pyrimidine Analogues on Geranyl Transferase. 2: 610 doi:10.4172/ V F L H Q W L ¿ F U H S R U W V

Copyright: © 2013 Ethiraj KR, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

from the binding site and redocking the six pyrimidines analogues

4, 6-trimethoxyphenyl) pyrimidin-2-amine>4-(2- uoro-4-methoxyphenyl)-6-(2,3, trimethoxyphenyl) pyrimidin-2-amine>4-(2- uoro-4-methoxyphenyl)-6-(2,4,5-trimethoxyphenyl) pyrimidin-2-amine> 4-(2- uoro-4-methoxyphenyl)-6-(2, 4-dimethoxyphenyl) pyrimidin-2-amine. Conformational analysis of di erent docked complexes also shows that residues Tyr 178, Ser 227 and Phe 230 play important role in this receptor's activity. Docking studies performed by GLIDE has con rmed that above inhibitors t into the binding pocket of the RabGGTase receptor. From the results, we may observe that for successful docking, intermolecular hydrogen bonding and liphophilic interactions between the ligand and the receptor are very important. We can also explain why the GLIDE score results for 4-(2- uoro-4-methoxyphenyl)-6-(3,4,5-trimethoxyphenyl) pyrimidin-2-amine are higher than others. e main reason for the increase in GLIDE score is due to the penalties for close intra-ligand contacts.

References

1. 2 N D P R W R & 7) R U W H - * 9 H V L F X O D U W U D I ¿ 3 N L Q H P D W L F D Q G L W V U H O D W L R Q V K L S W R F K R Q G U L D O F H O O (2009) Actin cytoskeleton, and H+-K+-ATPase recycling in the gastric parietal cell. J Physiol 532: 287-296.
2. Peter M, Chavrier P, Nigg EA, Zerial M (1992) Isoprenylation of rab proteins on structurally distinct cysteine motifs. J Cell Sci 102: 857-865.
3. Casey PJ, Seabra MC (1996) Protein prenyltransferases. J Biol Chem 271: 5289-5292.
4. Mattew J, Subba Rao AV, Rambhav S (1984) Propterol, an antibacterial agent from Pterocarpus marsupium. Curr Sci 53: 576-577.
5. Yamakawa T, Kagechika H, Kawachi E, Hashimoto Y, Shudo K (1990) Retinobenzoic acids. 5. Retinoidal activities of compounds having a trimethylsilyl or trimethylgermyl group(s) in human promyelocytic leukemia cells HL-60. J Med Chem 33: 1430-1437.
6. Bhat AK, Bhamana RP, Patel MR, Bellare RA, Deliwala CV (1972) Chemotherapy of fungus infections. III. Alkyl or aryl thiosemicarbazones, acid hydrazones, and styryl aryl ketones of 5-bromo- and 5-nitrosalicylaldehydes. Indian J Chem 10: 694-698.
7. Ishitsuka H, Ninomiya YT, Ohsawa C, Fujiu M, Suhara Y (1982) Direct and V S H F L ¿ F L Q D F W L Y D W L R Q R I U K L Q R Y L U X V E \ F K D O F Chemother 22: 617-621.
8. Ninomiya Y, Shimma N, Ishitsuka H (1990) Comparative studies on the antirhinovirus activity and the mode of action of the rhinovirus capsid binding agents, chalcone amides. Antiviral Res 13: 61-74.
9. Trivedi AR, Dodiya DK, Ravat NR, Shah VH (2008) Synthesis and biological evaluation of some new pyrimidines via a novel chalcone series. ARKIVOC 131-141.
10. Rahaman SkA, Rajendra Pasad Y, Phani Kumar, Bharath Kumar (2009) Synthesis and anti-histaminic activity of some novel pyrimidines. Saudi Pharmaceutical Journal 17: 255-258.
11. Jorgensen WL (2004) The many roles of computation in drug discovery. Science 303: 1813-1818.
12. Halgren TA, Murphy RB, Friesner RA, Beard HS, Frye LL, et al. (2004) Glide: a new approach for rapid, accurate docking and scoring. 2. Enrichment factors in database screening. J Med Chem 47: 1750-1759.
13. McRobb FM, Capuano B, Crosby IT, Chalmers DK, Yuriev E (2010) Homology modeling and docking evaluation of aminergic G protein-coupled receptors. J Chem Inf Model 50: 626-637.
14. Kaminski GA, Friesner RA (2001) Evaluation and Reparametrization of the